



ELSEVIER

# Targeting proteins to membranes: structure of the signal recognition particle

Pascal F Egea<sup>1</sup>, Robert M Stroud<sup>1</sup> and Peter Walter<sup>2</sup>

In all three kingdoms of life, co-translational targeting of secretory and membrane proteins to the prokaryotic plasma membrane or eukaryotic endoplasmic reticulum is mediated by a ribonucleoprotein complex, the signal recognition particle (SRP), and its membrane-associated receptor (SR). SRP binds to signal sequences of nascent proteins as they emerge from the exit tunnel of the ribosome. The resulting targeting complex, composed of the SRP and the ribosome–nascent chain complex (RNC), then docks with the SR in a GTP-dependent manner. Passing through a complex series of conformational states, SRP and SR deliver the RNC to the translocon, which in turn mediates protein translocation across or integration into the membrane. The core structural and mechanistic principles of SRP-dependent protein targeting are universally conserved. Recent structural investigations combining X-ray crystallography and cryo-electron microscopy have provided new insights into three essential steps of the SRP-dependent protein targeting cycle: the assembly and interaction of the SRP ribonucleoprotein core, the GTP-dependent SRP–SR association, and the interaction between SRP and the ribosome.

## Addresses

<sup>1</sup> Department of Biochemistry and Biophysics, University of California San Francisco, San Francisco, CA 94143, USA

<sup>2</sup> Howard Hughes Medical Institute, University of California San Francisco, San Francisco, CA 94143, USA

Corresponding author: Walter, Peter (pwalter@biochem.ucsf.edu)

**Current Opinion in Structural Biology** 2005, **15**:213–220

This review comes from a themed issue on  
Macromolecular assemblages  
Edited by Wolfgang Baumeister and Patrick Cramer

Available online 25th March 2005

0959-440X/\$ – see front matter  
© 2005 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.sbi.2005.03.007

## Introduction: the SRP-mediated targeting cycle

Evolutionarily related signal recognition particles (SRPs) and their cognate membrane-associated receptors (SRs) mediate the co-translational targeting of membrane and secretory proteins in all cells [1,2]. Signal sequences specify unidirectional protein translocation across or integration into membranes. A typical signal sequence consists of a stretch of 9–12 large hydrophobic residues that is either a transient or permanent part of the protein to be

translocated [3,4]. These sequences are likely to adopt  $\alpha$ -helical conformations when bound or in transit. If sufficiently hydrophobic but not too long, signal sequences can be recognized by the SRP as they emerge from ribosomes. Upon signal sequence recognition, SRP promotes a pause in translational elongation in eukaryotes, presumably to keep the nascent chain as short as possible before insertion into the translocation pore.

Both SRP and SR contain GTPase domains. Their signal-bound, GTP-dependent association into a tight complex is crucial for targeting the ribosome–nascent chain complex (RNC) to the protein translocation apparatus at the membrane. SRP–SR complex formation leads to reciprocal activation of GTP hydrolysis by both GTPases, which, only in complex, form a unique ‘dual active site’ in which the O3' of one GTP becomes a key component of the active site of the other GTPase. This reciprocity is thought to be coupled to and thus to govern docking, release of the RNC to the translocon and SRP recycling (Figure 1). Thus, SRP and SR act as molecular match-makers that deliver RNCs that synthesize selected subsets of proteins to the translocon. Recent structures and biochemical analysis provide new insights into the mechanism of SRP-mediated protein targeting [5,6].


This review gives an overview of the structural organization of the SRP throughout the diverse phyla, showing their evolutionary conservation, and focuses on the dynamic nature of the SRP-mediated protein targeting cycle. A recent cryo-EM structure of the eukaryotic SRP bound to the ribosome reveals an integrated picture of the interaction between SRP and the active ribosome. Also, the recent X-ray structure of the universally conserved heterodimeric core of the SRP–SR targeting complex reveals the crucial role of GTP for targeting complex assembly and unidirectionality of the targeting cycle. Combined with a recent structure of the protein translocation pore, these new insights raise fascinating questions about the molecular mechanisms of protein translocation through biological membranes.

## Architecture and assembly of SRPs in the different kingdoms of life: eukarya and archaea compared to bacteria

Metazoan SRPs consist of six proteins (SRP54, SRP19, SRP68, SRP72, SRP9, SRP14) and a 300-nucleotide RNA (SRP RNA) [7–9]. SRP can be divided into the Alu and S domains, which define the two major functional units of the SRP particle (Figure 2a). The S domain contains well-characterized SRP19 and SRP54, [10,11<sup>\*</sup>,12<sup>\*\*</sup>,13<sup>\*\*</sup>] and



Figure 2



The SRP and its receptor through evolution. Schematic representations of the architecture of the SRPs and SRs from (a) eukarya, (b) archaea and (c) bacteria. The GTPases (*Ffh*/*SRP54* and *FtsY*/*SRα*) are indicated in bold. The principal RNA helices present in eukaryotic and archaeal SRP RNA (~300 nucleotides), and in bacterial 4.5S RNA (~115 nucleotides) are labeled h2 through h8. In *SRP54*/*Ffh*, the M domain is responsible for signal peptide and RNA recognition. The NG domains of SRP and SR are closely related. Some bacterial *FtsY* contain an extra N-terminal A domain, which is proposed to be involved in association with the membrane. The eukaryotic SR is composed of two subunits: the regulatory subunit *SRβ* (containing an N-terminal transmembrane anchor) and *SRα*. *SRα* is subdivided into two domains: an N-terminal *SRx* domain involved in GTP-dependent *SRα*-*SRβ* heterodimerization and the NG domain. Some Gram-positive bacteria, such as *Bacillus subtilis*, retain a long SRP RNA with an Alu-like domain to which a dimeric protein (*HBsu*) is bound, akin to *SRP9*/*SRP14* in eukaryotic SRP. The main structural domains of SRP. (d) Structure of the human S domain, with the *SRP19* (PDB codes 1L1W, 1L9A and 1LNG) and *SRP54M* (PDB code 1MFQ) subunits bound, respectively, to helices h6 and h8 of the SRP RNA. (e) Structure of the human Alu domain (PDB codes 1914, 1E80 and 1E8S), with the *SRP9*/*SRP14* heterodimer bound to helices h3 and h4 of the SRP RNA. (f) Structure of the conserved NG domain of all SRP-type GTPases (PDB codes 1FTS, 1FFH, 1NG1, 2NG1, 3NG1, 1JPJ, 1JPN, 1O87 and 1J8M).

also contain an insertion box domain (IBD) that is unique to this GTPase subfamily (Figure 2f) [24–28].

The SRs are similarly phylogenetically conserved. In bacteria and archaea, SRs are single-subunit proteins, also named *FtsY* because the *E. coli* SR maps to an *Fts*

operon [29]. Bacterial SRs are peripherally associated with membranes (Figure 2b,c). By contrast, eukaryotic SRs are composed of two subunits: *SRα*, the homolog of *FtsY*, and *SRβ*, a membrane-anchoring subunit that contains yet another GTPase domain of unknown function (Figure 2a).

### Interaction with the translation machinery: ribosome and SRP

High-resolution X-ray structures of subcomplexes of eukaryotic, archaeal and bacterial SRPs, low-resolution EM structures of free SRP [30,31] and cryo-EM structures of SRP bound to RNCs [32\*\*] imply quite detailed structural and functional models for how the SRP-ribosome-SR interaction encodes correct targeting (Figure 3a). Comparison of the free and ribosome-bound mammalian SRP structures delineates major conformational changes and structural flexibility in SRP that bring it into its functionally critical state for targeting.

SRP binds to ribosomes whether or not a signal sequence is present; however, its affinity for ribosomes with emerging signal peptide is greater. Thus, SRP is thought to sample many RNCs efficiently for the presence of a signal sequence before identifying its correct 'signal'. Cross-linking localizes several anchoring points for SRP on the ribosome. Cross-links between residues of ribosomal proteins L23 and L35 (positioned in close proximity to the exit site of the protein tunnel through which the nascent chain traverses the large ribosomal subunit) and the N domain of SRP54 imply a zone of protein-protein contacts [33,34], and cross-links between SRP RNA and rRNA suggest SRP RNA-rRNA contacts [35]. Both are functionally important. The binding area for the Alu domain on the ribosome overlaps with the elongation factor binding site at the interface between the two ribosomal subunits [36], thus explaining how SRP modulates translation elongation (Figure 3b).

The structural basis of signal sequence binding and recognition remains mysterious. Several M domain structures have been solved [20,21,37,38] but, in spite of many attempts, no complex between SRP54 and a cognate signal sequence has yet been reported. The structures reveal that the M domain contains a deep groove that is bounded by a finger loop on one side and is lined by flexible sidechains of conserved hydrophobic residues (Figure 3c). The hydrophobic groove with structural plasticity is a conserved feature of M domains and may explain how SRP can bind signal peptides of diverse lengths and sequences. The conformational flexibility of the so-called 'finger loop' of the M domain may also play a role in the transition between the closed (empty) and open (signal-bound) states of the M domain. Signal sequences often contain positively charged residues that may play an as yet unrecognized role in SRP-signal peptide recognition through additional electrostatic interactions with the SRP RNA backbone.

Crystal structures of archaeal *Sulfolobus solfataricus* (*SsoI*) SRP54 in its free state and in complex with helix 8 of SRP RNA [39\*] revealed intrinsic flexibility between protein domains, and suggest that the hinge region linking the G and M domains (Figure 3d) is involved in interdomain

communication. Hydrophobic contacts between residues of the M and N domains could participate in communicating the occupancy state of the signal-binding pocket to the G domain. Biochemical data suggest that signal peptide binding might also involve the NG domain [40]. Thus, signal peptide binding could regulate the GTPase activity of SRP54 [41]. Analysis of the eukaryotic RNC-SRP and *SsoI* SRP structures further suggests that the flexible linker region plays a central role in a sequence of conformational changes [42]. It has been suggested that transfer of the signal sequence from the ribosome to the M domain results in an increased GTP affinity [43] that renders SRP competent for targeting of the RNC through specific interaction with the SR and subsequent docking of the RNC to the translocon. However, although this model is appealing on teleological grounds, at physiological concentrations, SRP is likely to be already occupied by GTP.

### Targeting to the membrane: the GTP catalytic cycle in protein targeting


Despite their similarities with conventional GTPases such as ras, SRP-type GTPases exhibit unique properties, including low affinity for, and rapid exchange of GDP and GTP. There is no known requirement for a guanine nucleotide exchange factor (GEF). Moreover, activation of SRP-type GTPases is triggered after SRP-SR complex formation between the two homologous GTPase domains of SRP54 and SR $\alpha$  [44]. In the SRP-SR complex, each GTPase acts as a GTPase-activating protein (GAP) for the other GTPase [45]; there is no known requirement for an extrinsic GAP.

How does SRP-SR direct correct targeting? Two recent X-ray structures [46\*\*,47\*\*] revealed structural insights into the mechanism of interaction of the GTPases in the SRP-SR complex. Both GTPases undergo conformational changes and associate as a head-to-head, quasi-twofold symmetrical heterodimer [48,49], burying an extensive interaction surface of 3200 Å<sup>2</sup>. Conserved residues from the N and G domains of both proteins contribute to the interface. A conserved ALLEADV sequence in the N domain and a conserved loop in the IBD define the edges of the heterodimer interface (see Figure 4a,b). The highly cooperative formation of the complex aligns the two GTP molecules in a shared composite active site, in which each GTP is contacted exclusively by sidechains contributed by the same GTPase to which it is bound.

The catalytically important interaction that spans the interface occurs between the two bound GTPs: the 3'-OH of one GTP is hydrogen bonded to the  $\gamma$ -phosphate of the other (Figure 4c) and vice versa. This circle of interactions between the twinned GTPs is severed twice upon hydrolysis, leading to complex dissociation after cargo delivery. Extensive biochemical analysis supports




Figure 3



The structure of mammalian SRP and its dynamic interactions with the ribosome. **(a)** Molecular model of the eukaryotic SRP based on the fitting of X-ray structures of eukaryotic, archaeal and bacterial SRP subcomplexes in the 12 Å resolution cryo-EM density map (PDB code 1RY1). Mouse and human SRP9/SRP14, *M. jannaschii* SRP19 and *M. jannaschii* SRP54 are displayed. The approximate location of the SRP68/SRP72 heterodimer is shown based on unassigned electron density in the cryo-EM maps. **(b)** Model of the eukaryotic SRP binding to an RNC. Free SRP is likely to adopt an extended conformation. SRP54 binds next to the peptide tunnel exit site and scans for signal peptides. Upon binding to the signal sequence emerging from the ribosome, SRP undergoes a conformational change (localized around the hinge 1 region near the SRP68/SRP72 heterodimer) and adopts the kinked conformation observed in the cryo-EM study. This promotes competitive binding of the Alu domain at the entry site for elongation factors (EF) and causes elongation arrest. **(c)** Signal peptide recognition by the M domain of SRP. The M domain of *T. aquaticus* Ffh has been superimposed on the structure of the *E. coli* ribonucleoprotein core (PDB codes 1DUL and 2FFH). The putative hydrophobic binding site for the signal peptide is shown (yellow). The finger loop adopts a variety of conformations in the several known structures and may play a role in the signal-binding process. **(d)** Conformational flexibility of SRP54/Ffh (PDB codes 2FFH, 1QZW and 1QZX). The structure of *SsoI* SRP54 bound to helix 8 of SRP RNA is shown with the N, G and M domains. The flexible linker between the G and M domains is shown in pink. **(e)** A model of the SRP conformational changes on the RNC. The exit tunnel of the 60S subunit of the ribosome is schematized with the reported cross-linking sites between SRP and ribosome. As the ribosome binds to SRP, it induces a conformational change that allows SRP to scan and sample for emerging signal peptides. Further conformational changes are transmitted to the GTPase core, resulting in SR binding. The association between the two NG domains partially exposes the translocon-binding site on the ribosome (brown arrow), initiating the docking of the RNC to the membrane.

Figure 4



The SRs and the universally conserved GTPase core of the SRP-SR targeting complex. **(a,b)** Structure of the *T. aquaticus* FtsY-Ffh NG complex (PDB codes 1OKK and 1RJ9). The two twinned SRP GTPases associate in a parallel way to form a quasi-symmetric heterodimer with an extensive surface that includes both N and G domains. Interaction of SRP (Ffh) with SR (FtsY) brings the two half active sites and their respective GTPs in close apposition to form the shared active site at the center of the two associated G domains. **(c)** Close-up view of the composite active site, with the twinned substrates, the essential and strictly conserved catalytic residues, the attacking waters (aw) and the magnesium cofactors. **(d)** Eukaryotic SR is composed of two subunits: SR $\alpha$  and SR $\beta$ . The structure of the yeast SR $\beta$ -SR $\alpha$  heterodimer (PDB code 1NRJ) shows the GTP-dependent interaction between the SR $\beta$  subunit (homologous to the Arf ras-like GTPases) and the SR $\alpha$  subunit. The NG domain of SR $\alpha$ , which is responsible for the interaction with the SRP, is not present in this structure. **(e)** A speculative eukaryotic SR cycle. GTP-primed SR $\beta$  binds the SR $\alpha$  domain of the SR $\alpha$  subunit. The SRP-RNC is targeted to the membrane via the GTP-dependent interaction between SRP and SR $\alpha$ , and docked on the translocon. It is unknown whether SR $\beta$  hydrolyzes GTP during each round of targeting and how this hydrolysis is regulated. The translocon might function as a nucleotide exchange factor for SR $\beta$ .

the importance and functional significance of the extensive interaction surface, and the role of the 3'-OH groups in association, reciprocal activation and catalysis [46<sup>••</sup>,50<sup>•</sup>]. The IBD motifs rearrange and each contribute three strictly conserved residues, which are of central importance to catalysis. Two of these residues contact the  $\gamma$ -phosphate and stabilize the transition state, whereas the catalytic aspartate residue positions a water molecule for nucleophilic attack, which will then break the  $\beta$ - $\gamma$  phosphate bond (Figure 4c). It is possible that an additional arginine residue has to be repositioned to complete the active site [47<sup>••</sup>,50<sup>•</sup>].

These structures provide the first insight into how the SR-SRP interaction drives SRP-dependent protein targeting. The structures explain the coupling of SRP-SR complex formation and its subsequent disassembly, and suggest a unique activation mechanism for the SRP family of GTPases. The extensive interactions, both from active site residues and from the twinned substrates,

explain why complex formation is GTP dependent and how GTP hydrolysis leads to complex dissociation.

Distinct classes of mutations have been isolated that inhibit specific steps of SRP-SR complex formation and activation, suggesting an ordered series of discrete conformational steps during formation of the active SRP-SR complex [6,50<sup>•</sup>]. Each step could potentially provide a control point in the targeting reaction by the action of additional components, thus ensuring the binding and release of cargo at the appropriate place and time.

In eukaryotes, an additional level of complexity arises from the heterodimeric structure of SR, in which the additional GTPase SR $\beta$  is an essential subunit. The association between the two subunits, SR $\alpha$  and SR $\beta$ , is GTP dependent, as shown by the recent structure of the yeast SR $\alpha$ -SR $\beta$  heterodimer [51<sup>••</sup>] (Figure 4d). SR $\beta$  has been suggested to coordinate signal sequence release from SRP with ribosome binding to the translocon

[52,53] (Figure 4e). As such, SR $\beta$  might represent an additional layer of regulation in the transfer of signal sequences from the SRP to the translocon. Alternatively, SR $\beta$  may use its GTPase domain to control recruitment and activation of SR $\alpha$  in a way that more globally controls the translocation capacity of the cell according to need.

## Conclusions

Several essential questions concerning SRP-dependent protein targeting remain to be answered. How is signal sequence recognition by SRP accomplished? We have not yet obtained a structure of such a complex. Furthermore, SRP RNA and signal sequence regulate different steps of the GTP catalytic cycle; SRP RNA stimulates the SRP–SR interaction [54] and dissociation of the signal sequence from the SRP appears to be a prerequisite to GTP hydrolysis [55]. Comparison of the SRP–RNC and RNC–translocon cryo-EM structures shows that both SRP and the translocon bind next to the exit site of the ribosomal protein tunnel [32<sup>\*\*</sup>,42,56,57] in a mutually exclusive way, indicating that targeting must coordinate SRP release with translocon binding. The mechanisms by which SRP facilitates the transfer of the nascent chain and the docking of the ribosome onto the translocon remain unknown [58,59<sup>\*\*</sup>]. The structural flexibility of SRP appears to be crucial for it to act as an adaptor between the ribosome and the translocon.

## Acknowledgements

This work was supported by grants from the National Institutes of Health to RMS (GM060641), PW and PFE, and funds from The Herbert Boyer Fund to PFE. PW is an investigator of the Howard Hughes Medical Institute.

## References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Keenan RJ, Freymann DM, Stroud RM, Walter P: **The signal recognition particle**. *Annu Rev Biochem* 2001, **70**:755-775.
  2. Eichler J, Moll R: **The signal recognition particle of archaea**. *Trends Microbiol* 2001, **9**:130-136.
  3. von Heijne G: **Targeting sequences**. In *Protein Targeting, Transport & Translocation*. Edited by Dalbey RE, von Heijne G. Academic Press; 2002:35-46.
  4. Stroud RM, Walter P: **Signal sequence recognition and protein targeting**. *Curr Opin Struct Biol* 1999, **9**:754-759.
  5. Doudna JA, Batey RT: **Structural insights into the signal recognition particle**. *Annu Rev Biochem* 2004, **73**:539-557.
  6. Shan SO, Walter P: **Co-translational protein targeting by the signal recognition particle**. *FEBS Lett* 2005, **579**:921-926.
  7. Nagai K, Oubridge C, Kuglstatter A, Menichelli E, Isel C, Jovine L: **Structure, function and evolution of the signal recognition particle**. *EMBO J* 2003, **22**:3479-3485.
  8. Wild K, Weichenrieder O, Strub K, Sinning I, Cusack S: **Towards the structure of the mammalian signal recognition particle**. *Curr Opin Struct Biol* 2002, **12**:72-81.
  9. Sauer-Ericksson AE, Hainzl T: **S-domain assembly of the signal recognition particle**. *Curr Opin Struct Biol* 2003, **13**:64-70.
  10. Wild K, Sinning I, Cusack S: **Crystal structure of an early protein-RNA assembly complex of the signal recognition particle**. *Science* 2001, **294**:598-601.
  11. Oubridge C, Kuglstatter A, Jovine L, Nagai K: **Structure of SRP19 in complex with the S domain of SRP RNA and its implication for the assembly of the signal recognition particle**. *Mol Cell* 2002, **9**:1251-1261.  
The authors describe the crystal structure of a non-cognate binary complex between archaeal *Methanococcus jannaschii* SRP19 and the human S domain of 7S RNA.
  12. Hainzl T, Huang S, Sauer-Ericksson AE: **Structure of the SRP19 RNA complex and implications for signal recognition particle assembly**. *Nature* 2002, **417**:767-771.  
The crystal structure of the archaeal binary complex formed between SRP19 and the S domain of the 7S RNA from *M. jannaschii* is reported. It illustrates the molecular basis of SRP19 binding to the S domain and the overall architecture of the RNA, showing the stabilizing interactions established between RNA helices 6 and 8, which stack against each other to form the S domain.
  13. Kuglstatter A, Oubridge C, Nagai K: **Induced structural changes of 7SL RNA during the assembly of human signal recognition particle**. *Nat Struct Biol* 2002, **9**:740-744.  
This article presents the X-ray structure of the ternary complex between human SRP19, SRP54 (M domain) and the S domain of 7S RNA. Binding of the M domain induces conformational changes in the RNA. A mechanism of S domain assembly in humans is proposed.
  14. Birse DE, Kapp U, Strub K, Cusack S, Åberg A: **The crystal structure of the signal recognition particle *Alu* RNA binding heterodimer SRP9/14**. *EMBO J* 1997, **16**:3757-3766.
  15. Weichenrieder O, Wild K, Strub K, Cusack S: **Structure and assembly of the *Alu* domain of the signal recognition particle**. *Nature* 2000, **408**:167-173.
  16. Powers T, Walter P: **Co-translational protein targeting catalyzed by the *Escherichia coli* signal recognition particle and its receptor**. *EMBO J* 1997, **16**:4880-4886.
  17. Rose MA, Weeks KM: **Visualizing induced fit in early assembly of the human signal recognition particle**. *Nat Struct Biol* 2001, **8**:515-520.
  18. Maeshima H, Okuno E, Aimi T, Morinaga T, Itoh T: **An archaeal protein homolog to mammalian SRP54 and bacterial Ffh recognizes a highly conserved region of SRP RNA**. *FEBS Lett* 2001, **507**:336-340.
  19. Tozik I, Huang Q, Zwieb C, Eichler J: **Reconstitution of the signal recognition particle of the halophilic archaeon *Haloflex volcanii***. *Nucleic Acids Res* 2002, **30**:4166-4175.
  20. Batey RT, Sagar MB, Doudna J: **Structural and energetic analysis of RNA recognition by a universally conserved protein from the signal recognition particle**. *J Mol Biol* 2001, **307**:229-246.
  21. Batey RT, Doudna J: **Structural and energetic analysis of metal ions essential to SRP signal recognition domain assembly**. *Biochemistry* 2002, **41**:11703-11710.
  22. Keenan RJ, Freymann DJ, Walter P, Stroud RM: **Crystal structure of the signal sequence binding subunit of the signal recognition particle**. *Cell* 1998, **94**:181-191.
  23. Freymann DM, Keenan RJ, Stroud RM, Walter P: **Structure of the conserved GTPase domain of the signal recognition particle**. *Nature* 1997, **385**:361-364.
  24. Freymann DM, Keenan RJ, Stroud RM, Walter P: **Functional changes in the structures of the SRP GTPase on binding GDP and Mg<sup>2+</sup> GDP**. *Nat Struct Biol* 1999, **6**:793-801.
  25. Montoya G, Te Kaat K, Moll R, Schafer G, Sinning I: **The crystal structure of the conserved GTPase core from the archaeon *Acidianus ambivalens* and its comparison with related structures suggests a model for SRP-SRP receptor complex**. *Structure* 2000, **8**:515-525.
  26. Padmanabhan S, Freymann DM: **The conformation of bound GMPNP suggests a mechanism for gating the active site of the SRP GTPase**. *Structure* 2001, **9**:859-867.



27. Ramirez U, Minasov G, Focia PJ, Stroud RM, Walter P, Juhn P, Freymann DM: **Structural basis for mobility in the 1.1 Å crystal structure of the NG domain of *Thermus aquaticus* Ffh.** *J Mol Biol* 2002, **320**:783-799.
28. Focia PJ, Alam H, Lu T, Ramirez U, Freymann DM: **Novel protein and Mg<sup>2+</sup> configurations in the Mg<sup>2+</sup> GDP complex of the SRP GTPase Ffh.** *Proteins* 2004, **54**:222-230.
29. Montoya G, Svensson C, Lührink J, Sinning I: **Crystal structure of the NG domain from the signal recognition particle receptor FtsY.** *Nature* 1997, **385**:365-368.
30. Andrews DW, Walter P, Ottensmeyer FP: **Structure of the signal recognition particle by electron microscopy.** *Proc Natl Acad Sci USA* 1985, **82**:785-789.
31. Andrews DW, Walter P, Ottensmeyer FP: **Evidence for an extended 7SL RNA structure in the signal recognition particle.** *EMBO J* 1987, **6**:3471-3477.
32. Halic M, Becker T, Pool MR, Spahn CMT, Grassucci RA, Frank J, Beckmann R: **Structure of the signal recognition particle interacting with the elongation arrested ribosome.** *Nature* 2004, **427**:808-814.
- The authors present cryo-EM reconstructions of SRP-RNC complexes. By fitting the known X-ray structures of SRP and SRP RNA subdomains in the cryo-EM density maps, the authors propose a model of both the interaction between SRP, the ribosome and a nascent chain, and SRP-mediated translation elongation arrest.
33. Gu S, Peske F, Wieden H, Rodnina MV, Wintermeyer W: **The signal recognition particle binds to protein L23 at the peptide exit of the *Escherichia coli* ribosome.** *RNA* 2003, **9**:566-573.
34. Pool MR, Stumm J, Fulga TA, Sinning I, Dobberstein B: **Distinct modes of signal recognition particle interaction with the ribosome.** *Science* 2002, **297**:1345-1348.
35. Rinke-Appel J, Osswald M, von Knoblauch K, Mueller F, Brimacombe R, Sergiev P, Adveeda O, Bogdanov A, Dontsova O: **Crosslinking of 4.5S RNA to the *Escherichia coli* ribosome in the presence or absence of the protein Ffh.** *RNA* 2002, **8**:612-625.
36. Terzi L, Pool MR, Dobberstein B, Strub K: **Signal recognition particle Alu domain occupies a defined site at the ribosomal subunit interface upon signal sequence recognition.** *Biochemistry* 2004, **43**:107-117.
37. Batey RT, Rambo RP, Lucast L, Rha B, Doudna J: **Crystal structure of the ribonucleoprotein core of the signal recognition particle.** *Science* 2000, **287**:1232-1239.
38. Clemons WM Jr, Gowda K, Black SD, Zwieb C, Ramakrishnan V: **Crystal structure of the conserved subdomain of human protein SRP54M at 2.1 Å resolution: evidence for the mechanism of signal peptide binding.** *J Mol Biol* 1999, **292**:697-705.
39. Rosendal KR, Wild K, Montoya G, Sinning I: **Crystal structure of the complete core of archaeal signal recognition particle and implications for inter-domain communication.** *Proc Natl Acad Sci USA* 2003, **100**:14701-14706.
- This article reports the low-resolution crystallographic structure of the SRP from the archaeon *SsoI* in the absence or presence of a fragment of the SRP RNA. This is the first structure of a full-length SRP54 in which the linker between the NG and M domains is observed.
40. Cleverley RM, Gierasch LM: **Mapping the signal sequence-binding site on SRP reveals a significant role for the NG domain.** *J Biol Chem* 2002, **277**:46763-46768.
41. Wild K, Rosendal KR, Sinning I: **A structural step into the SRP cycle.** *Mol Microbiol* 2004, **53**:357-363.
42. Wild K, Halic M, Sinning I, Beckmann R: **SRP meets the ribosome.** *Nat Struct Mol Biol* 2004, **11**:1049-1053.
43. Bacher G, Lutcke H, Jungnickel B, Rapoport TA, Dobberstein B: **Regulation by the ribosome of the GTPase of the signal recognition particle during protein targeting.** *Nature* 1996, **381**:248-251.
44. Mandon EC, Gilmore R: **GTPase twins in the SRP family.** *Nat Struct Mol Biol* 2004, **11**:115-116.
45. Powers T, Walter P: **Reciprocal stimulation of GTP hydrolysis by two directly interacting GTPases.** *Science* 1995, **269**:1422-1424.
46. Egea PF, Shan S-O, Napetschnig J, Savage DF, Walter P, Stroud RM: **Substrate twinning activates the signal recognition particle and its receptor.** *Nature* 2004, **427**:215-221.
- This paper (and [47\*\*]) is the first to report the high-resolution structure of the universally conserved heterodimeric core of a targeting complex comprising the twinned GTPase domains of an SRP (Ffh) and its cognate SR (FtsY) from *Thermus aquaticus*. This structure reveals the crucial role of GTP in SRP-SR association and reciprocal activation, thus explaining the unidirectionality of the targeting reaction.
47. Focia PJ, Shepotinovskaya IV, Seidler JA, Freymann DM: **Heterodimeric GTPase core of the SRP targeting complex.** *Science* 2004, **303**:373-377.
- See annotation to [46\*\*].
48. Jagath JR, Rodnina MV, Wintermeyer W: **Conformational changes in the bacterial SRP receptor FtsY upon binding of guanine nucleotides and SRP.** *J Mol Biol* 2000, **295**:745-753.
49. Shepotinovskaya IV, Freymann DM: **Conformational change of the N-domain on formation of the complex between the GTPase domains of *Thermus aquaticus* Ffh and FtsY.** *Biochim Biophys Acta* 2002, **1597**:107-114.
50. Shan S-O, Stroud RM, Walter P: **Mechanism of association and reciprocal activation of two GTPases.** *PLoS Biol* 2004, **2**:e320.
- This article presents an extensive biochemical analysis of the FtsY-Ffh interaction, combining site-directed mutagenesis and kinetics. It dissects the various stages of bacterial SRP-SR complex formation and activation.
51. Schwartz T, Blobel G: **Structural basis for the function of the β subunit of the eukaryotic signal recognition particle.** *Cell* 2003, **112**:793-803.
- This article presents the first X-ray structure of a complex between the regulatory domain (SR $\alpha$ ) of eukaryotic SR $\alpha$  and its membrane-docking subunit, SR $\beta$ .
52. Fulga TA, Sinning I, Dobberstein B, Pool MR: **SR $\beta$  coordinates signal sequence release from SRP with ribosome binding to the translocon.** *EMBO J* 2001, **20**:2338-2347.
53. Mandon EC, Jiang Y, Gilmore R: **Dual recognition of the ribosome and the signal recognition particle by the SRP receptor during protein targeting to the endoplasmic reticulum.** *J Cell Biol* 2003, **162**:575-585.
54. Peluso P, Hershlag D, Nock S, Freymann D, Johnson AE, Walter P: **Role of 4.5S RNA in assembly of the bacterial signal recognition particle with its receptor.** *Science* 2000, **288**:1640-1643.
55. Song W, Maden D, Gilmore R: **Role of Sec61 $\alpha$  in the regulated transfer of the ribosome-nascent chain complex from the signal recognition particle to the translocation channel.** *Cell* 2000, **100**:333-343.
56. Beckmann R, Spahn CM, Eswar N, Helmers J, Penczek PA, Sali A, Frank J, Blobel G: **Architecture of the protein conducting channel associated with the translating 80S ribosome.** *Cell* 2001, **107**:361-372.
57. Morgan DG, Ménétret JF, Neuhof A, Rapoport TA, Akey CW: **Structure of the mammalian ribosome-channel complex at 17 Å resolution.** *J Mol Biol* 2002, **324**:871-886.
58. Breyton C, Haase W, Rapoport TA, Kühlbrandt W, Collinson I: **Three-dimensional structure of the bacterial protein-translocation complex SecYEG.** *Nature* 2002, **418**:662-665.
59. Van den Berg B, Clemons WM Jr, Collinson I, Modis Y, Hartmann E, Harrison SC, Rapoport TA: **X-ray structure of a protein conducting channel.** *Nature* 2003, **427**:36-44.
- The authors report the first crystallographic structure of a translocon (SecYEG complex), from the archaeon *M. jannaschii*.