1	Engineering ER-stress dependent non-conventional mRNA splicing
2	
3	Weihan Li ^{1, *} , Voytek Okreglak ^{1, 3} , Jirka Peschek ¹ , Philipp Kimmig ^{1, 4} , Meghan
4	Zubradt ² , Jonathan Weissman ² , Peter Walter ^{1, *}
5	
6	¹ Department of Biochemistry and Biophysics and Howard Hughes Medical
7	Institute, University of California San Francisco, CA 94143, USA
8	² Department of Cellular and Molecular Pharmacology and Howard Hughes
9	Medical Institute, University of California San Francisco, CA 94143, USA
10	³ Current address: Calico Life Sciences LLC, South San Francisco, CA 94080, USA
11	⁴ Current address: Institute of Biochemistry, ETH Zürich, Zürich, Switzerland
12	
13	*: Corresponding authors
14	

16 Abstract

17 The endoplasmic reticulum (ER) protein folding capacity is balanced with the 18 protein folding burden to prevent accumulation of un- or misfolded proteins. The ER 19 membrane-resident kinase/RNase Ire1 maintains ER protein homeostasis through 20 two fundamentally distinct processes. First, Ire1 can initiate a transcriptional 21 response through a non-conventional mRNA splicing reaction to increase the ER 22 folding capacity. Second, Ire1 can decrease the ER folding burden through selective 23 mRNA decay. In Saccharomyces cerevisiae and Schizosaccharomyces pombe, the two 24 Ire1 functions have been evolutionarily separated. Here, we show that the 25 respective Ire1 orthologs have become specialized for their functional outputs by 26 divergence of their RNase specificities. In addition, RNA structural features separate 27 the splicing substrates from the decay substrates. Using these insights, we 28 engineered an *S. pombe* Ire1 cleavage substrate into a splicing substrate, which 29 confers *S. pombe* with both Ire1 functional outputs. 30

31 Introduction

33	In eukaryotes, the vast majority of secretory and transmembrane proteins
34	are folded in the endoplasmic reticulum (ER). The ER protein folding homeostasis is
35	maintained by a collective of signaling pathways, termed the unfolded protein
36	response (UPR) (Walter and Ron, 2011, Ron and Walter, 2007). The most
37	evolutionarily conserved branch of the UPR is mediated by the ER-transmembrane
38	kinase/endoribonuclease (RNase) Ire1. Direct binding of unfolded proteins to Ire1's
39	ER lumenal domain triggers Ire1 to oligomerize and form foci (Gardner and Walter,
40	2011, Karagoz et al., 2017, Credle et al., 2005, Aragon, van Anken et al., 2009). In
41	turn, Ire1 oligomerization activates Ire1's cytosolic kinase/RNase domain
42	(Korennykh et al., 2009), which restores ER homeostasis through two functional
43	outputs. First, Ire1 initiates a process of non-conventional cytosolic splicing of <i>XBP1</i>
44	mRNA (in metazoans) or HAC1 mRNA (in S. cerevisiae). Translation of the spliced
45	mRNA produces a transcription factor Xbp1 (Hac1 in <i>S. cerevisiae</i>), which drives a
46	large transcriptional program to adjust the ER's protein-folding capacity according
47	to the protein folding load in the ER lumen (Cox et al., 1993, Mori et al., 1993,
48	Yoshida et al., 2001, Calfon et al., 2002, Sidrauski et al., 1996). Second, Ire1 can
49	reduce the ER folding burden by cleaving a set of mRNAs encoding ER-target
50	proteins. The initial Ire1-mediated cleavage leads to mRNA degradation, in a
51	process termed regulated Ire1-dependent decay (RIDD) (Hollien and Weissman,
52	2006, Hollien et al., 2009, Kimmig, Diaz et al., 2012). The mechanism that

distinguishes the non-conventional mRNA splicing from RIDD has largely remainedunknown.

55

56 Interestingly, the two Ire1 modalities co-exist in metazoan cells (Hollien and 57 Weissman, 2006, Hollien et al., 2009, Moore and Hollien, 2015), yet are 58 evolutionarily separated in the two yeast species, S. cerevisiae and S. pombe. The 59 UPR in *S. cerevisiae* engages Ire1 exclusively in mRNA splicing, whereas in *S. pombe* 60 it engages Ire1 exclusively in RIDD. There is no detectable RIDD in *S. cerevisiae* and 61 no *HAC1* / *XBP1* ortholog in *S. pombe*, nor is there a corresponding transcriptional 62 program (Niwa et al., 2005, Kimmig, Diaz et al., 2012). It is intriguing to note that the 63 fundamental task of maintaining ER protein homeostasis can be achieved by two radically different processes catalyzed by two distantly related Ire1 orthologs. The 64 65 two yeast species, *S. cerevisiae* and *S. pombe*, therefore provide a unique opportunity 66 to dissect the two Ire1 functional outputs, which has remained an unsolved 67 challenge in metazoans. Here, we set out to exploit this opportunity.

- 69 Results
- 70

71 S. pombe and S. cerevisiae Ire1's cytosolic domains are functionally divergent

72

73 Despite their common role as UPR effectors and conserved domain structure 74 (Fig. 1A), S. pombe and S. cerevisiae Ire1 orthologs share 29% sequence identity, and 75 sequence variation may confer Ire1's functional divergence. To explore this notion, 76 we swapped the homologous *IRE1* genes between the two yeast species by 77 expressing *S. cerevisiae* Ire1 in *Δire1 S. pombe* cells and, *vice versa*, *S. pombe* Ire1 in 78 *Aire1 S. cerevisiae* cells. To this end, we constructed strains in which we integrated 79 the foreign IRE1 genes into the genomes of the other yeast such that their 80 expression was regulated by the host species-endogenous IRE1 promoters and the 81 resulting mRNAs contained host species-endogenous 5' and 3' untranslated regions 82 (UTR). The IRE1 genes contained sequences encoding FLAG-tags that we inserted 83 into an unstructured loop in their ER-lumenal domains in a position known to 84 preserve Ire1 function (Rubio et al., 2011). In both yeasts, the foreign genes 85 expressed Ire1 at comparable levels (Fig. 1B & C, lanes 3). However, when grown on plates containing tunicamycin, a drug that blocks N-linked glycosylation and induces 86 87 ER stress, the foreign Ire1s failed to support cell growth of either *S. pombe* and *S.* 88 *cerevisiae* cells (Fig. 1D & E, lanes 3), indicating that *S. pombe* and *S. cerevisiae* Ire1s 89 are not interchangeable.

91 There are two plausible, not mutually exclusive scenarios that could explain 92 the failure of cross-species complementation: i) the foreign Ire1 lumenal domains 93 fail to sense ER stress, or ii) the foreign Ire1 cytosolic domains fail to recognize 94 species-appropriate RNA substrates. Since the Ire1 lumenal domains have lower 95 sequence identity (21%) than the cytosolic kinase/RNase domains (45%), we first 96 swapped the Ire1 lumenal domains, generating chimeras with foreign lumenal 97 domains and host species-endogenous transmembrane/cytosolic domains. Both 98 chimeras supported growth on tunicamycin plates, suggesting that the divergent 99 Ire1 lumenal domains share a conserved mechanism to sense ER stress and 100 transduce the signal across ER membrane (Fig. 1D & E, lanes 4). Next, we swapped 101 the Ire1 transmembrane/cytosolic domains. These Ire1 chimeras failed to restore 102 growth on tunicamycin plates of both S. pombe and S. cerevisiae cells (Fig. 1D & E, 103 lanes 5), indicating that the Ire1 transmembrane/cytosolic domains cause the Ire1 104 functional incompatibility when expressed in the opposing yeast. As a control, we 105 expressed FLAG-tagged host species-endogenous Ire1s into Δ ire1 strains of both 106 yeasts. These strains phenocopied the wild type (WT) cells on tunicamycin plates 107 (Fig. 1D & E, lanes 6). We again confirmed by immunoblotting that all of the FLAG-108 tagged Ire1 constructs were stably expressed at near-endogenous level in both 109 yeasts (Fig. 1B & C).

110

111 We next asked whether the Ire1 constructs would process the host species-112 appropriate RNA substrates in *S. pombe* and *S. cerevisiae* cells. To this end, we 113 performed Northern blot and qPCR analyses to measure cleavage and subsequent

114 down-regulation of *GAS2* mRNA, which is a RIDD target in *S. pombe* cells (Kimmig, Diaz et al., 2012). We performed the Northern blot in $\Delta ski2$ *S. pombe*, in which the 115 116 RNA 3' to 5' decay machinery is impaired so that the *GAS2* mRNA 5' cleavage 117 fragments can be detected in the gel. Of the different Ire1 variants, only the Ire1 118 chimera bearing the *S. pombe* cytosolic domain cleaved the *GAS2* mRNA (Fig. 1— 119 figure supplement 1A) and decreased the mRNA level (Fig. 1F), consistent with the 120 growth phenotype. In S. pombe, Ire1 also cleaves the BIP1 mRNA within its 3'UTR, 121 producing a truncated mRNA with an increased half-life (Kimmig, Diaz et al., 2012). 122 To assess *BIP1* mRNA processing, we performed qPCR analysis using two pairs of 123 primers, one pair bracketing the Ire1 cleavage site and the other pair bracketing a 124 region upstream of it (Fig. 1G, schematic insert, *black* vs. *grev* arrows), reporting on 125 uncleaved only and both *BIP1* mRNA species (i.e. total *BIP1* mRNA), respectively. As 126 expected, upon tunicamycin-induced ER stress in WT cells uncleaved BIP1 mRNA 127 levels decreased while total BIP1 mRNA level increased (Fig. 1G, lanes 1, 2). As 128 shown in Figure 1G, Ire1 variants bearing the *S. pombe* cytosolic domain processed 129 *BIP1* mRNA, whereas Ire1 variants bearing the *S. cerevisiae* cytosolic domain did not. 130 This result is further validated by Northern blot analysis of *BIP1* mRNA (Fig. 1— 131 figure supplement 1B).

132

Correspondingly in *S. cerevisiae* cells, we examined *HAC1* mRNA splicing by
PCR across its splice junction. Consistent with the cell growth phenotype on
tunicamycin, the two Ire1 constructs bearing the *S. pombe* cytosolic domains did not
splice *HAC1* mRNA in *S. cerevisiae* cells (Fig. 1H, lanes 6, 10). By contrast, the Ire1

137	chimera bearing the <i>S. pombe</i> lumenal domain and <i>S. cerevisiae</i> cytosolic domains		
138	spliced <i>HAC1</i> mRNA (Fig. 1H, lane 8), albeit at reduced efficiency compared to WT <i>S</i> .		
139	cerevisiae Ire1 (Fig. 1H, lane 2). We confirmed the activity of the various Ire1		
140	constructs in HAC1 mRNA splicing by monitoring UPR dynamics with a HAC1 mRNA-		
141	derived splicing reporter (Fig. 1—figure supplement 2A & B) previously described		
142	(Aragon, van Anken et al., 2009, Zuleta et al., 2014). The reduced HAC1 mRNA		
143	splicing efficiency observed for Ire1 bearing the S. pombe lumenal domain (Fig. 1H,		
144	lane 8, and Fig. 1—figure supplement 2B, column 4) can be explained by the		
145	observation that the <i>S. pombe</i> lumenal domain mediates a lower degree of		
146	oligomerization than its <i>S. cerevisiae</i> counterpart, as demonstrated by the reduced		
147	ability of Ire1-mCherry fusion constructs to form foci visible by fluorescent		
148	microscopy (Fig. 1—figure supplement 2C). Consistent with previous studies		
149	(Aragon, van Anken et al., 2009), the insertion of the mCherry module into the Ire1		
150	cytosolic linker, which connects Ire1 transmembrane domain and cytosolic		
151	kinase/RNase domain, did not affect its ability to sustain cell growth (Fig. 1—figure		
152	supplement 2D).		
153			
154	Ire1 kinase/RNase domains have distinct RNase specificity		
155			
156	To further confine the Ire1 region giving rise of the species differences in		
157	outputs, we expressed an Ire1 chimera that, in addition to the <i>S. cerevisiae</i> lumenal		
158	domain, also included the S. cerevisiae transmembrane and cytosolic linker domains		
159	fused to the <i>S. pombe</i> kinase/RNase domain. This chimeric Ire1 weakly rescued cell		

growth and mildly restored the *HAC1* mRNA splicing upon ER stress (Fig. 2A lane 4,
& Fig. 2B lane 6), compared to the chimera containing *S. pombe* transmembrane and
cytosolic linker domains, although both constructs were expressed at similar
protein levels (Fig. 2C). This result indicates that the major difference lies in the
kinase/RNase domains, but that the transmembrane and cytosolic linker domains
can afford a marginal rescue, most likely by reintroducing cytosolic linker elements
that facilitate *HAC1* mRNA docking (van Anken et al., 2014).

167

168 To study the differences by which the Ire1 kinase/RNase domains select 169 their respective substrate mRNAs, we purified recombinant S. cerevisiae and S. 170 pombe kinase/RNase domains and performed in vitro RNA cleavage assays. The S. 171 cerevisiae Ire1 kinase/RNase efficiently cleaved a cognate 29-nucleotide RNA 172 hairpin derived from the 3' splice site of *S. cerevisiae HAC1* mRNA (Fig. 2D). By 173 contrast, under the same conditions the S. pombe Ire1 kinase/RNase cleaved the S. 174 cerevisiae HAC1 mRNA-derived substrate ~60-fold slower (Fig. 2D). Reciprocally, 175 the *S. pombe* Ire1 kinase/RNase efficiently cleaved a cognate 32-nucleotide RNA 176 hairpin derived from the cleavage site in *S. pombe BIP1* mRNA and cognate RNA 177 hairpins derived from the RIDD cleavage sites in *S. pombe SPAC4G9.15* and *PLB1*, 178 whereas S. cerevisiae Ire1 kinase/RNase cleaved the BIP1 mRNA-derived substrate 179 and the *S. pombe SPAC4G9.15* >500-fold slower and *PLB1* mRNA-derived hairpins 180 ~100-fold slower (Fig. 2E, F, G). These in vitro data validate and expand the 181 conclusions from the experiments conducted *in vivo*, suggesting that the different

182 Ire1 RNase specificities separate their functional outputs and that they are not183 dependent on other cellular factors such as associated proteins or lipids.

184

185 Ire1 kinase/RNase domains recognize distinct RNA sequence and structural 186 features

187

188 Ire1 recognizes its substrates based on both RNA sequence and structural 189 features. The required RNA sequence motifs were characterized previously and 190 differ between species: for *S. cerevisiae* Ire1 a nucleotide sequence of CNG/CNGN or 191 CNG|ANGN ("|" indicates the Ire1 cleavage position) situated in a strictly conserved 192 7-membered loop is required (Gonzalez et al., 1999). By contrast, for *S. pombe* Ire1 a 193 three-nucleotide sequence of UG|C is required and no additional structural features 194 have yet been characterized (Kimmig, Diaz et al., 2012, Guydosh et al., 2017). To fill 195 this gap in our knowledge, we examined the RNA secondary structures in vivo. To 196 this end, we treated the *S. pombe* cells with dimethyl sulfate (DMS), which allows 197 detection of exposed (unpaired and not blocked by proteins) adenine/cytosine 198 residues (illustrated as green dots in Fig. 3A). RNA was extracted, reverse 199 transcribed and deep-sequenced. The DMS modifications stop reverse transcriptase 200 and generate truncated DNA fragments that we mapped through deep sequencing. 201 We then used identified unpaired bases to guide *in silico* RNA secondary structure 202 predictions (Rouskin et al., 2014). For example, near one of the GAS2 mRNA 203 cleavage sites, five bases, labeled in green in Figure 3B, have high DMS modification 204 signals. In the RNA folding software mfold (Zuker, 2003), we provided the

205 constraint such that these five residues are unpaired and obtained the predicted 206 RNA secondary structure (Fig. 3C). In this structure, the GAS2 mRNA forms a 9-207 membered stem loop with the cleavage consensus sequence UG|C located near the 208 center of the loop. In similar analyses of 13 additional *S. pombe* Ire1 substrate mRNA 209 cleavage sites previously identified by both Kimmig, Diaz et al. and Guydosh et al. 210 (Guydosh et al., 2017, Kimmig, Diaz et al., 2012), we found in all of them cleavage 211 sites located near the center of loops in RNA stem-loop structures (Fig. 3—figure 212 supplement 1 A, B). By contrast to those found in S. cerevisiae HAC1 mRNA, the 213 predicted loops were of variable sizes, with the smallest being a 3-membered loop 214 (e.g., the SPAC4G9.15 mRNA cleavage site) and the largest being a 9-membered loop 215 (e.g., the *BIP1* mRNA cleavage site). We summarize that *S. pombe* Ire1 is tolerant to 216 loop size variation, while the *S. cerevisiae* Ire1 stringently recognizes 7-membered 217 stem loops. Thus, the S. cerevisiae and S. pombe Ire1 recognize distinct RNA 218 sequence and structural features (Fig. 3D). 219 220 A prediction of this model is that RNAs that combine *S. cerevisiae* and *S.* 221 *pombe* Ire1 motifs should be substrates to Ire1 from either species. To test this 222 prediction, we analyzed a substrate satisfying criteria for both species. Specifically, 223 we examined a substrate predicted to form a 7-membered stem loop with the 224 sequence CUG|CAGC, meeting the criteria of both the *S. cerevisiae* Ire1 motif 225 (CNG|CNGN) and the *S. pombe* Ire1 motif (UG|C). Indeed, both enzymes cleaved this 226 RNA *in vitro* with similar efficiency, in strong support of our model (Fig. 3E).

227

228	We further challenged the model <i>in vivo</i> by modifying the <i>S. pombe BIP1</i>
229	mRNA. First, we replaced the Ire1 cleavage site in the 3' UTR of <i>BIP1</i> mRNA with a
230	sequence derived from the S. cerevisiae HAC1 mRNA 5' splice site. This modification
231	is predicted to change the endogenous 9-membered loop to a 7-membered one. The
232	new construct (' <i>BIP1-HAC1</i> hybrid mRNA') contained the sequence AG C, which is
233	different from the <i>S. pombe</i> Ire1 sequence motif UG C. In support of our model, <i>S.</i>
234	pombe Ire1 failed to cleave the BIP1-HAC1 hybrid mRNA upon UPR induction (Fig.
235	3G, bar 3 & 4, Fig. 3—figure supplement 2A, B). Next, we mutated the non-cognate A
236	to a cognate U, such that the <i>S. pombe</i> Ire1 cleavage motif UG C was restored. Indeed,
237	the single nucleotide change restored the <i>S. pombe</i> Ire1 cleavage (Fig. 3G, bar 5 & 6,
238	Fig. 3—figure supplement 2C). Taken together, we conclude that the Ire1 orthologs
239	in <i>S. cerevisiae</i> and <i>S. pombe</i> have divergent substrate preferences.
240	
241	Engineering non-conventional mRNA splicing in <i>S. pombe</i>
242	
243	While these results revealed the differences between the <i>S. cerevisiae</i> and <i>S.</i>
244	pombe UPR at the step of Ire1 cleavage, it was not clear what determines the fates of
245	the RNA cleavage fragments, i.e. RNA ligation in S. cerevisiae and RNA degradation in
246	S. pombe. To address this question, we tested whether S. pombe cells contain
247	functional mRNA ligation machinery. To this end, we expressed the S. cerevisiae
248	<i>HAC1</i> mRNA-derived splicing reporter (Fig. 1—figure supplement 2A) in Δ <i>ire1 S.</i>
249	pombe cells, bearing genomic copies of various Ire1 constructs. A chimeric Ire1
250	bearing the <i>S. pombe</i> cytosolic domain failed to splice the reporter mRNA, in

agreement with our model (Fig. 4A, lane 5 & 6). Interestingly, both Ire1 constructs
bearing the *S. cerevisiae* cytosolic domains successfully spliced the reporter mRNA
(Fig. 4A, lanes 1-4). Thus, the *S. pombe* cells ligated the mRNA cleavage fragments as
long as the correct RNA substrates were provided. This result suggested that
features in the RNA substrates determine their fate post Ire1 cleavage.

256

257 Recently, we reported that in mammalian cells the *XBP1* mRNA actively 258 participates in the splicing reaction. In particular, a conformational RNA 259 rearrangement promotes XBP1 mRNA intron ejection and exon ligation (Peschek, 260 Acosta-Alvear et al., 2015). We wondered if this mechanism could be the factor that 261 diverges the fates of the RNA cleavage fragments. To address this question, we 262 aimed to synthetically create the Ire1-dependent non-conventional mRNA splicing 263 reaction in *S. pombe* cells initiated by endogenous *S. pombe* Ire1. First, we identified 264 the analogous RNA conformational rearrangement in S. cerevisiae HAC1 mRNA. To 265 obtain a *HAC1*-derived RNA splicing cassette optimized for *S. pombe* Ire1, we then 266 engineered the two Ire1-cleavage sites at the splice junctions to match the *S. pombe* 267 Ire1 UG|C motif and pruned the intron (originally 252 bp in *S. cerevisiae*) to the very 268 residues predicted to be critical for the mRNA conformational rearrangement (30 269 bp). Finally, we inserted the S. pombe-optimized mRNA splicing cassette into S. 270 *pombe BIP1* mRNA, replacing its endogenous Ire1 cleavage site (Fig. 4B, Fig. 4–figure 271 supplement 1). Indeed, we found that the *BIP1* mRNA containing the synthetic 272 splicing cassette was spliced in *S. pombe* upon induction of ER stress (Fig. 4C lane 2). 273 Sequencing of the lower band in Figure 4C (lane 2) verified the designed identity of

274 the splicing product (Fig. 4D). To show that insertion of the splicing cassette triggers 275 mRNA splicing independent of particular flanking elements, we inserted the splicing 276 cassette into the 3'UTR of another synthetic mRNA. In this case, we constructed a 277 synthetic mRNA containing the 5' UTR of tubulin (*NDA2*) mRNA, the open reading 278 frame of a GAS2 mutant mRNA in which all of its RIDD cleavage sites were mutated, 279 and the 3' UTR of *NDA2* mRNA with the inserted splicing cassette (Fig. 4E). As for 280 the mRNA described above, this construct was efficiently spliced in *S. pombe* upon 281 ER stress (Fig. 4F). Thus, we conclude that the substrate RNA structure determines 282 the fate of the RNA cleavage fragments.

283

285 Discussion

286

287 "What I cannot create, I do not understand" (Richard Feynman). Inspired by 288 this quote, we engineered *S. pombe* cells to carry out a non-conventional mRNA 289 splicing reaction that does not occur in this species naturally. This feat was enabled 290 by a combination of a detailed mechanistic characterization of the differences in 291 Ire1-dependent mRNA processing between two yeast species reported in this paper 292 and by functional insights gleaned previously from a characterization of the 293 mammalian *XBP1* mRNA splicing mechanism (Peschek, Acosta-Alvear et al., 2015). 294 Two main conclusions emerged from this study. First, the evolutionary divergence 295 of Ire1's substrate specificity determines the set of mRNAs that are cleaved in either 296 species. Second, features in the mRNAs determine the distinct fates of the severed 297 mRNA fragments. S. pombe can be coopted to carry out the core non-conventional 298 mRNA splicing reaction with fidelity, as long as it is provided with a spliceable RNA 299 substrate that matches the specificity requirements of its endogenous Ire1. Albeit 300 not optimized in evolution for efficiency of the reaction, no other components (such 301 as the RNA ligase) need to be fundamentally specialized to mediate the core splicing 302 reaction.

303

304 Ire1 orthologs in *S. cerevisiae* and *S. pombe* recognize their cognate mRNA
305 substrates by discriminating both sequence and structural features. *S. pombe* Ire1
306 RNase specificity is more promiscuous, and has a broad substrate scope, in line with
307 its role to initiate degradation of many ER-bound mRNAs to reduce the ER's protein

308 folding burden. By contrast, *S. cerevisiae* Ire1's RNase specificity is very stringent 309 and specialized with HAC1 mRNA being as its only substrate in the cell (Niwa et al., 310 2005), in line with its role to produce a single transcription activator to drive UPR 311 target genes. In mammals, two paralogs of Ire1 are expressed in a tissue-specific 312 manner (Tsuru et al., 2013, Bertolotti et al., 2001). Ire1α, which performs both *XBP1* 313 mRNA splicing and RIDD, recognizes a similar but longer RNA sequence motif, 314 CUG/CAG, displayed in stem-loop structures (Maurel et al., 2014). Interestingly, the 315 loop sizes that Ire1α recognizes differ between the *XBP1* mRNA cleavage sites and 316 RIDD cleavage sites, with the two cleavage sites on XBP1 mRNA to be conserved 7-317 mer loops (Hooks and Griffiths-Jones, 2011), while the cleavage sites on RIDD 318 substrates vary in range from 9-mers to 5-mers (Fig. 4—figure supplement 2). This 319 suggests that mammalian Ire1 α may display two distinct modes of RNase activity— 320 a more stringent mode of RNase activity observed on 7-mer stem-loop RNAs (XBP1 321 mRNA and BLOC1S mRNA) and a more promiscuous mode of RNase activity on RNA 322 substrates with variable loop sizes. As shown here, these two modes have been 323 cleanly separated in evolution between *S. cerevisiae* and *S. pombe* Ire1. Hence it 324 seems plausible that mammalian Ire1 α may switch selectively into one or the other 325 state, perhaps in response to the timing of UPR activation or certain physiological 326 conditions, which could be reflected in particular oligomeric states, post-327 translational modifications, or other effectors yet to be discovered. 328 329 Ire1 α is more efficient in *XBP1* mRNA splicing, while Ire1 β prefers to cleave

ribosomal RNA (Iwawaki et al., 2001, Nakamura et al., 2011, Imagawa et al., 2008).

331 Pairwise sequence alignment did not reveal an obvious similarity signatures that 332 would distinguish the Ire1 species performing for RIDD, i.e., S. pombe Ire1 and Ire1β, 333 from those engaged in mRNA splicing, i.e., *S. cerevisiae* Ire1 and Ire1α (Fig. 4—figure 334 supplement 3A, B). On the substrate side, two Ire1 β cleavage sites both located on 335 rRNA, have been mapped to date. They share a common sequence of G|C at the 336 cleavage site. Previous studies indicated that differences in Ire1 α 's and Ire1 β 's 337 RNase domains lead to their functional distinction (Imagawa et al., 2008). We did 338 not observed cleavage of rRNA by *S. pombe* Ire1 (Fig. 1—figure supplement 1A, B); 339 mammalian Ire1 β 's activity to do so may therefore reflect a specialization that is not 340 generalizable to all Ire1 RNases that perform RIDD. Therefore, modulating Ire1's 341 RNase specificity to regulate its mode of action emerges as a general theme for 342 different species, as well as for different tissues within the same species.

343

344 In S. cerevisiae cells, apart from the HAC1 mRNA, other mRNAs contain the S. 345 cerevisiae Ire1 cleavage motif, yet are not cleaved (Niwa et al., 2005). This is 346 explained by spatial coordination. *HAC1* mRNA is targeted to Ire1 upon stress, 347 utilizing a specific signal in the HAC1 mRNA 3'UTR (Aragon, van Anken et al., 2009), 348 conferring exquisite specificity that renders *HAC1* mRNA the sole substrate of the 349 reaction. Although two other Ire1 substrate RNAs have been reported (Tam et al., 350 2014), we have not been able to reproduce this result. By contrast to the dedicated 351 mRNA targeting in S. cerevisiae, S. pombe and mammalian cells target XBP1 mRNA 352 and most RIDD substrates to the ER via the signal recognition particle (SRP) 353 pathway (Hollien and Weissman, 2006, Hollien et al., 2009, Yanagitani et al., 2009,

Yanagitani et al., 2011, Plumb et al., 2015). In this way, *S. pombe* and mammalian
Ire1 efficiently sample through substrate RNAs at ER periphery, which they cleave
more promiscuously. We presume that in our experimental set-up, the chimeric *BIP1*-mRNA containing the splicing cassette would highjack the SRP-mediated
targeting route initiated by the BiP1 signal sequence, as we previously showed for
other RIDD substrate mRNAs (Kimmig, Diaz et al., 2012).

360

361 After Ire1 cleavage, RNA fragments are ligated or degraded, depending on the 362 substrate RNA structures. According to this notion, in mammalian cells where 363 splicing and RIDD co-exist, we predict that *XBP1* mRNA splicing and RIDD substrate 364 degradation are separated post Ire1 cleavage. The two cleavage sites on the XBP1 365 mRNA are coordinated by a zipper-like RNA structure, which enable the exons to be 366 held in juxtaposition and ligated by the cytosolic tRNA ligase (Peschek, Acosta-367 Alvear et al., 2015, Sidrauski et al., 1996, Jurkin et al., 2014, Lu et al., 2014, 368 Kosmaczewski et al., 2014). By contrast, cleavage sites on the RIDD substrates lack 369 such coordination and the cleavage fragments are further degraded. 370

371Our study revealed that Ire1's RNase specificity and its RNA substrate372structure separate Ire1's modes of action, opening the door to identify residues that373shape Ire1's RNase specificity. In this way, it should now become possible to design374metazoan Ire1s that favor mRNA splicing over RIDD, and *vice versa*, enabling us to375discriminate the biological significance of the two Ire1 functional outputs separately376in physiological and pathological contexts.

377 Acknowledgement

379	The Δ <i>ire1 S. pombe</i> strain was a generous gift from the Krogan lab (UCSF). We
380	thank Ignacio Zuleta, Benjamin Heineike and Alain Bonny for their help and
381	discussion on automated flow cytometry. We thank Wallace Marshall, Marc Shuman
382	and members of the Walter lab for their insightful discussions. This work was
383	supported by UCSF-Zaffaroni Fellowship (WL), the Human Frontier Science
384	Program (JP, PK), National Science Foundation grant 1144247 (MZ), the Genentech
385	Foundation (MZ) and CRSB (Center for RNA Systems Biology; grant P50 GM102706
386	to JSW). PW and JSW are Investigators of the Howard Hughes Medical Institute.
387	
388	Competing Financial Interests
389	
390	The authors declare no competing financial interests.

Figure 1

393	S. pombe and S. cerevisiae Ire1 have functionally conserved stress sensing ER-
394	lumenal domains and divergent cytosolic domains. (A) Cartoon illustration of
395	lumenal domain (LD), transmembrane/cytosolic linker domain (TMD+L) and
396	kinase/RNase domain (KR) for <i>S. pombe</i> (<i>Sp</i>) (<i>blue</i>) and <i>S. cerevisiae</i> (<i>Sc</i>) Ire1
397	(orange). (B, C) Expression levels of S. cerevisiae Ire1 (128 kD), S. cerevisiae lumenal
398	S. pombe cytosolic Ire1 (126 kD), S. pombe lumenal S. cerevisiae cytosolic Ire1 (125
399	kD) and <i>S. pombe</i> Ire1 (122 kD) in <i>S. pombe</i> (B) and <i>S. cerevisiae</i> cells (C). Extracts
400	were immunoblotted for 3xFLAG-Ire1. Ponceau stain (B) or Pgk1 (C) was used as
401	loading control. (D, E) Cell growth assay on tunicamycin (Tm) plates. Serial dilutions
402	of <i>S. pombe</i> (D) or <i>S. cerevisiae</i> (E) cells, which expressed the indicated Ire1
403	constructs, were spotted onto plates containing 0.05 μ g/ml (B) or 0.1 μ g/ml (C) of
404	Tm. Plates were photographed after incubation at 30° C for 4 days. (F, G) qPCR assay
405	for S. pombe GAS2 (F) or BIP1 (G) mRNA fold change upon 1 μ g/ml Tm treatment for
406	1 h. Experiments were done in triplicates. In (G), uncleaved (<i>dark grey</i>) or total
407	(<i>light grey</i>) <i>BIP1</i> mRNA was detected using the corresponding PCR primers
408	illustrated as arrows in the schematic insert. The red dashed line indicates the Ire1
409	cleavage position on <i>BIP1</i> mRNA. (H) Detection of <i>S. cerevisiae HAC1</i> mRNA splicing
410	by RT-PCR across the splice junction. Cells were treated with or without 1 $\mu g/ml$ of
411	Tm for 1 h.
412	

- **Figure 1—figure supplement 1**
- 414 Ire1 chimeras with *S. pombe* cytosolic domain cleave *BIP1* and *GAS2* mRNA in *S.*
- *pombe*. Northern blots of *S. pombe GAS2* (A) and *BIP1* (B) mRNA. Cells were treated
- 416 with $1 \mu g/ml$ of Tm for 1 h.

418 Figure 1—figure supplement 2

419 Ire1 oligomeric state determines the *HAC1* mRNA splicing dynamics in *S.*

420 *cerevisiae* cells. (A) Illustration of the *HAC1* mRNA derived splicing reporter. The

421 splicing reporter contains a part of the *HAC1* mRNA 5' exon replaced by the green

422 fluorescent protein (GFP) coding sequence. In the unspliced reporter mRNA,

423 translation is inhibited by a translation block formed by the intron and 5'UTR

424 (indicated by the red arrow). Upon splicing, intron is removed and translation

425 begins. (B) Measuring the *HAC1* mRNA splicing dynamics in *S. cerevisiae* using

426 automated flow cytometry. After 1.5 h of incubation, either no Tm or 0.25 μ g/ml, 0.5

427 μ g/ml, 1 μ g/ml, 2 μ g/ml of Tm was added. Then, we monitored the splicing

428 dynamics for 10 h. The splicing dynamics under various conditions is plotted. *Green*

429 lines represent the strains of interest, which expressed indicated Ire1 variants, and

430 the *black* line represents WT control strain under the same condition. (C)

431 Examining Ire1 foci formation in *S. cerevisiae* cells via fluorescence microscopy with

432 or without 1 μg/ml Tm treatment for 20 min. (D) Growth assay on Tm plate for *S*.

433 *cerevisiae* cells expressing Ire1 constructs with or without mCherry inserted into the

434 cytosolic linker. The inserted mCherry does not affect Ire1's ability to alleviate ER

435 stress.

436

437 Figure 2

438 S. pombe and S. cerevisiae Ire1 have distinct RNase specificity. (A) Growth assay 439 for *S. cerevisiae* cells expressing indicated Ire1 constructs on Tm plates, as Figure 1E. 440 (B) Measuring *HAC1* mRNA splicing, as Figure 1H. (C) Comparing the expression 441 levels of the indicated 3xFLAG-tagged Ire1 chimeras using immunoblotting. Ponceau 442 stain was used as loading control. (D, E, F, G) In vitro RNA cleavage assays. 5'-443 radiolabeled hairpin RNA substrates were incubated with 12.5 µM S. cerevisiae or S. 444 *pombe* Ire1 kinase/RNase domains (KR) at 30° C for the indicated time. (D) Hairpin 445 RNA substrate derived from the 3' splice site of S. cerevisiae HAC1 mRNA. The calculated k_{obs} is 9.4 \pm 0.9 \times 10 $^{\text{-4}}$ s $^{\text{-1}}$ for S. cerevisiae Ire1 KR and 0.15 \pm 0.01 \times 10 $^{\text{-4}}$ s $^{\text{-1}}$ 446 447 for *S. pombe* Ire1 KR. (E) Hairpin RNA substrate derived from the Ire1 cleavage site 448 on *S. pombe BIP1* mRNA. The calculated k_{obs} is 0.079 ± 0.0006 × 10⁻⁴ s⁻¹ for *S.* 449 *cerevisiae* Ire1 KR and 37.3 \pm 4.4 \times 10⁻⁴ s⁻¹ for *S. pombe* Ire1 KR. (F) Hairpin RNA 450 substrate derived from the Ire1 cleavage site on S. pombe SPAC4G9.15 mRNA, 451 encoding a gene of unknown function. The calculated k_{obs} was below our detection 452 limit for S. cerevisiae Ire1 KR and $15.6 \pm 2.2 \times 10^{-4}$ s⁻¹ for S. pombe Ire1 KR. (G) 453 Hairpin RNA substrate derived from the Ire1 cleavage site on *S. pombe PLB1* mRNA. 454 The calculated k_{obs} is 0.2 ± 0.003 × 10⁻⁴ s⁻¹ for *S. cerevisiae* Ire1 KR and 19.0 ± 2.5 × 455 10⁻⁴ s⁻¹ for *S. pombe* Ire1 KR.

457 **Figure 3**

458 S. pombe and S. cerevisiae Ire1 recognize distinct RNA sequence and structural 459 features. (A) Illustration of RNA structural mapping by DMS modifications. 460 Dimethyl sulfate (DMS) allows detection of unpaired adenine and cytosine RNA 461 bases (green dots). (B) The normalized DMS modification signals near the Ire1 462 cleavage site on *S. pombe GAS2* mRNA (cleavage site is indicated by the *red* dashed 463 line). The positions with high DMS modification signals are labeled in *green* and the 464 previously identified S. pombe Ire1 UG|C motif is labeled in red. (C) In sillico RNA 465 secondary structure prediction of the Ire1 cleavage site on GAS2 mRNA. Structure 466 prediction was constrained by forcing the positions with high DMS modification 467 signals (green) to be unpaired. (D) RNA sequence and structural motifs recognized 468 by the *S. cerevisiae* and *S. pombe* Ire1. (E) *In vitro* cleavage assay using an RNA 469 hairpin derived from human XBP1 mRNA 3' splice site, which is predicted to be a 470 shared substrate for *S. cerevisiae* and *S. pombe* Ire1 KR. The calculated kobs is 16.7 ± 471 2.3×10^{-4} s⁻¹ for *S. cerevisiae* Ire1 KR and $38.9 \pm 4.0 \times 10^{-4}$ s⁻¹ for *S. pombe* Ire1 KR. 472 (F) Illustrations of the *S. pombe BIP1* mRNA variants and (G) their uncleaved (*dark* 473 *grey*) or total (*light grey*) mRNA fold change upon ER stress in *S. pombe* cells. 474 Experiments were done in triplicates.

- 476 Figure 3—figure supplement 1
- 477 *S. pombe* Ire1 cleaves at UG|C positioned near the center of loops in RNA stem-
- 478 **loop structures.** (A) A list of all 14 *S. pombe* Ire1 mRNA cleavage sites, which were
- 479 independently identified by both Kimmig, Diaz et al. and Guydosh et al. (Guydosh et
- 480 al., 2017, Kimmig, Diaz et al., 2012). The UG|C motifs are labeled in *red*. The
- 481 positions with high DMS modification signals are labeled in green. (B) Predicted
- 482 RNA secondary structures of *S. pombe* Ire1 cleavage sites. DMS modification signals
- 483 were used to guide the secondary structure prediction of *S. pombe* Ire1 mRNA
- 484 cleavage sites. The *red* dashed lines indicate the Ire1 cleavage sites.

- 486 Figure 3—figure supplement 2
- 487 Ire1 cleavage sites on *BIP1* mRNA variants. Sequence and predicted RNA
- 488 secondary structures of Ire1 cleavage sites on (A) S. pombe BIP1 mRNA, (B) BIP1-
- 489 HAC1 hybrid mRNA and (C) BIP1-HAC1 hybrid mRNA with an A to U mutation. The
- 490 part included in the dashed box is derived from *S. cerevisiae HAC1* mRNA 5' splice
- 491 site.
- 492

493 **Figure 4**

494 Engineering the Ire1-mediated non-conventional mRNA splicing in *S. pombe*

- 495 **cells.** (A) Measuring the non-conventional mRNA splicing in *S. pombe* cells, which
- 496 were transformed with the *S. cerevisiae HAC1* mRNA splicing reporter and the
- 497 indicated Ire1 constructs. Cells were treated with $1 \mu g/ml$ Tm for 1 h. (B)
- 498 Illustration of the engineered *S. pombe BIP1* mRNA splicing variant. (C) Measuring
- the nonconventional mRNA splicing of the engineered *S. pombe BIP1* mRNA splicing
- 500 variant. Experimental conditions are the same as those for Figure 4A. (D)
- 501 Sequencing reads of the spliced *BIP1* mRNA. The schematic illustrations (E) and the
- 502 splicing assays (F) of the synthetic splicing substrates in *S. pombe*. Cells were treated
- 503 with 1 μ g/ml Tm for 1 h.

Figure 4—figure supplement 1

The splicing cassette in the engineered *S. pombe BIP1* **mRNA splicing variant**.

- 507 The part included in the dashed box is the inserted synthetic splicing cassette. The
- *red* dashed lines indicate the Ire1 cleavage sites. The *S. pombe* Ire1 UG|C motifs are
- 509 labeled in red.

Figure 4—figure supplement 2

The Ire1 *α* **cleavage sites on** *XBP1* **mRNA and a RIDD targets.** Red dashed lines

- 513 mark the cleavage sites and the red letters indicate the previously identified
- 514 sequence motif.

- 516 **Figure 4—figure supplement 3**
- 517 The sequence alignment of the kinase/RNase domains of Ire1 α , Ire1 β , the *S*.
- 518 *cerevisiae* Ire1 and the *S. pombe* Ire1. (A) The sequence alignment and colored
- 519 with BoxShade Server. (B) The sequence identities between the indicated pairs of
- 520 Ire1 constructs.
- 521

522 Materials and Methods:

523

524 Strains, plasmids and growth conditions

525 Standard *S. cerevisiae* and *S. pombe* genome editing and growth conditions were

- 526 used (Moreno et al., 1991, Guthrie and Fink, 2002). Strains used in this study are
- 527 listed in the Table 1. Specifically, all Ire1 constructs have a 3x FLAG-tag in their
- 528 lumenal domains replacing an unstructured region (255-274 in S. pombe and 267-
- 529 286 in *S. pombe*). *S. cerevisiae* Ire1 domain boundaries were previously described
- 530 (Rubio et al., 2011), *S. pombe* Ire1 domains were determined by sequence alignment
- with *S. cerevisiae* Ire1. Specifically, the lumenal domain is 1-526 for *S. cerevisiae* and

532 1-507 for *S. pombe*. The transmembrane/cytosolic linker is 527-672 for *S. cerevisiae*

- 533 and 508-651 for *S. pombe*. Kinase/RNase is 673-1115 for *S. cerevisiae* and 652-1073
- for *S. pombe*. Ire1 constructs were integrated into the *HO* locus in *S. cerevisiae*
- 535 (backbone plasmid: HO-Poly-KanMX4-HO) and *Leu* locus in *S. pombe* (backbone
- plasmid: pJK148). *S. pombe BIP1* variants were integrated at the *BIP1* locus through
- 537 homologous recombination and uracil selection. The mCherry-tagged Ire1
- 538 constructs and the splicing reporter were previously described (Aragon, van Anken
- 539 et al., 2009).
- 540

541 Growth assay

- 542 Serial dilutions of *S. cerevisiae* or *S. pombe* cells were spotted onto YPD plates with
- 543 0.1 μg/ml tunicamycin (for *S. cerevisiae*) or YE5S plates with 0.05 μg/ml

tunicamycin (for *S. pombe*). Plates were photographed after incubating at 30° C for 4days.

546

547 Immunoblotting

548 For both *S. cerevisiae* and *S. pombe* cells, total protein was isolated from yeast

549 cultures growing at exponential phase by vortexing with glass beads in 8 M urea, 50

550 mM Hepes, pH 7.4, and 1% sodium dodecylsulfate (SDS). Samples were boiled and

then centrifuged at 16,000 x g for 10 min. A sample containing 20 μ g total protein

552 was separated using electrophoresis and then transferred to nitrocellulose. The

553 3xFLAG-tagged Ire1 was probed with monoclonal anti-FLAG antibody (Sigma

554 F3165).

555

556 **qPCR assays**

557 Total RNA was purified from yeast cultures using phenol extraction (Kohrer and

558 Domdey, 1991). RNA samples were resuspended in RNase-free water and quantified

by spectrophotometry. cDNA was synthesized by reverse transcription using

560 random hexamer DNA primers (Thermo Fisher Scientific), SuperScript II Reverse

561 Transcriptase (Thermo Fisher Scientific) and 1 µg total RNA as described previously

562 (Kimmig, Diaz et al., 2012). 1% of the cDNAs was employed for qPCR reactions using

563 SYBR green qPCR kit (Bio-Rad). qPCR was performed in triplicates using CFX96

564 Touch Real-Time PCR Detection System (Bio-rad). qPCR primers are listed in Table

565 2. mRNA levels were normalized to *NDA2* mRNA in *S. pombe*.

566

567 In vivo mRNA splicing assay

- 568 cDNA was synthesized the same way as described in the qPCR section. Then we
- 569 used Phusion High-Fidelity PCR Kit (NEB) and performed PCR with cDNA and a set
- 570 of primers across the splice junction. For *HAC1* mRNA, the forward primer was
- 571 ATGGAAATGACTGATTTTGAACTAACTAGTAATTCG. The reverse primer was
- 572 TCATGAAGTGATGAAGAAATCATTCAAATTCAAATG. The PCR was performed for 26
- 573 cycles with annealing temperature of 51.5° C and extension time of 30 s. For the *S*.
- 574 *pombe BIP1* mRNA containing the splicing cassette, the forward primer was
- 575 GAATCGTGACTCTATAGCCATTAACA. The reverse primer was
- 576 CAATTATTGTCAGTTCCACAAAGC. The PCR was performed for 36 cycles with
- annealing temperature of 63.4° C and extension time of 15 s. For *S. cerevisiae HAC1*
- 578 mRNA derived splicing reporter expressed in *S. pombe* cells, the forward primer was
- 579 GAACTACAAGACACGTGCTGAAG. The reverse primer was
- 580 GATGAAGAAATCATTCAAATG. The PCR was performed for 60 cycles with
- annealing temperature of 63.2° C and extension time of 20 s. For the synthetic
- 582 splicing substrate in *S. pombe*, the forward primer was
- 583 CTCATTTAGATTAGCAATTCAAATG. The reverse primer was
- 584 GATTAGATCAACAATTCAAATGATC. The PCR was performed for 40 cycles with
- annealing temperature of 59.7° C and extension time of 20 s.

586

587 **Recombinant protein purification**

- 588 *S. cerevisiae* Ire1 kinase/RNase was purified as previously described (Korennykh et
- al., 2009). Details of the *S. pombe* Ire1 kinase/RNase purification will be described in

590 a separate paper. Briefly, *S. pombe* Ire1 kinase/RNase was N-terminally fused with 591 Glutathione S-transferase (GST) tag through a linker containing Human Rhinovirus 592 (HRV) 3C protease cleavage site, and was regulated by T7 promoter. This S. pombe 593 Ire1 kinase/RNase expression cassette was transformed into *E. coli* cells. 16 h after 594 transformation, we mixed and collected all the colonies on the transformation plates 595 by scraping them off from the agar plate into 50 ml of LB medium. After 3-hour 596 incubation at 37° C, the sample was diluted to 12 l of LB medium and further 597 incubated at 37° C until optical density reached 1. Protein expression was induced 598 by adding 0.5 mM IPTG. Then, the culture was incubated at 25° C for 4 h before we 599 pelleted the cells by centrifugation. Cells were resuspended in GST binding buffer 600 (50 mM Tris/HCl pH 7.5, 500 mM NaCl, 2 mM Mg(OAc)₂, 2 mM DTT, 10% Glycerol) 601 and homogenized using high-pressure homogenizer (EmulsiFlex, Avestin). The cell 602 lysate was applied to GST-affinity column and eluted with GST elusion buffer (50 603 mM Tris/HCl pH 7.5, 200 mM NaCl, 2 mM Mg(OAc)₂, 2 mM DTT, 10% Glycerol, 10 604 mM glutathione). The column elution was treated with GST-tagged HRV 3C protease 605 (PreScission Protease, GE Health). At the same time, the sample was dialyzed to 606 remove glutathione in the elution buffer. Next, the sample was further purified 607 through negative chromatography by passing through a GST-affinity column (to 608 remove free GST and residue GST-fused Ire1 kinase/RNase) and an anion exchange 609 column (to remove contaminating nucleic acids). Finally, the sample was subject to 610 gel filtration, concentrated to about 14 µM in storage buffer (50 mM Tris/HCl pH 611 7.5, 200 mM NaCl, 2 mM Mg(OAc)₂, 2 mM TCEP, 10% Glycerol), and flash frozen in

612 liquid nitrogen. The final purity, as well as purity at intermediate steps, was

613 assessed by SDS-PAGE using Coomassie blue staining.

614

615 In vitro RNA cleavage assays

616 Short RNA oligos were purchased from Dharmacon, Inc. RNA oligos were gel 617 extracted, acetone precipitated and resuspended in RNase-free water. Then, oligos 618 were 5' end radio-labeled with γ -[³²P]-ATP (Perkin Elmer) using T4 polynucleotide 619 kinase (NEB) and cleaned using ssDNA/RNA Clean and Concentrator kit (Zymo 620 Research D7010). To fold the RNA oligos, we heated the RNA oligos to 90° C for 3 621 min and slowly cooled them down at a rate of 1° C per minute until the temperature 622 reached 10° C. In the Ire1 cleavage assays, the reaction samples contained 12.5 µM 623 of *S. cerevisiae* or *S. pombe* Ire1 kinase/RNase. The cleavage reaction was performed 624 at 30° C in reaction buffer (50 mM Tris/HCl pH 7.5, 200 mM NaCl, 2 mM Mg(OAc)₂, 2 625 mM TCEP, 10% Glycerol). At each time point, an aliquot of 0.75 µl was transferred to 626 5 µl STOP buffer (10 M urea, 0.1% SDS, 1 mM EDTA, 0.05% xylene cyanol, 0.05% 627 bromophenol blue). RNAs were separated using denaturing 15% urea-PAGE gels 628 (run at 100 V for 90 min). Gels were imaged with a Phosphorimager (Typhoon FLA 629 9500, GE Health) and the band intensities were quantified using image. The cleaved 630 portion was calculated as the cleaved band intensity divided by the sum of the 631 cleaved band and uncleaved band intensities. The kobs were obtained by fitting the 632 data to first-order ("one-phase") decay equation using Prism. For the cleavage 633 reactions that less than 10% of the substrates were cleaved, because the substrate concentration was approximately constant, the cleavage dynamics was fitted to a 634

- 635 linear equation to obtain k_{obs}. The sequence of hairpin RNA substrate derived from
- 636 Ire1 cleavage site on *S. pombe BIP1* mRNA is
- 637 CGCGAGAUAACUGGUGCUUUGUUAUCUCGCG.
- 638 The sequence of hairpin RNA substrate derived from Ire1 cleavage site on *S. pombe*
- 639 *SPAC4G9.15* mRNA is CCACCACCGAGUAUGCUACUCGGUGGUGG.
- 640 The sequence of hairpin RNA substrate derived from *S. cerevisiae HAC1* mRNA 3'
- 641 splice site is GCGCGGACUGUCCGAAGCGCAGUCCGCGC
- 642 The sequence of hairpin RNA substrate derived from *XBP1* mRNA 3' splice site is
- 643 UGCACCUCUGCAGCAGGUGCA.
- 644

645 Automated flow cytometry

- 646 Measuring *S. cerevisiae* UPR dynamics using automated flow cytometry was
- 647 previously described in detail (Zuleta et al.). Briefly, we co-cultured two *S. cerevisiae*
- 648 strains, a strain of interest and a control strain. The control strain contained a
- 649 constitutively expressed mCherry reporter. The signal from the control strain was
- 650 computationally separate based on its high mCherry level. In an 11.5-hour time
- 651 course at 30° C, a data point was taken every 20 min. 1.5 h after inoculation, DMSO
- 652 (as control) or 0.25 μg/ml, 0.5 μg/ml, 1 μg/ml, 2 μg/ml of Tm were added. Splicing
- 653 dynamics were monitored for another 10 h. The GFP signal was normalized to the
- 654 signal at time zero.

655

656 Probing in vivo mRNA structure in S. pombe cells

657 A culture of 15 ml *S. pombe* cells, which were exponentially growing at 30° C, was

658 treated with 400 µl of DMS for 3.5 min. DMS was then quenched by adding 30 ml of

659 solution containing 30% β-mercaptoethanol and 25% isoamyl alcohol. Then, cells

660 were pelleted by centrifugation at 4° C, and washed with 15 ml 30% β -

- 661 mercaptoethanol. Total RNA was extracted using phenol extraction. Poly(A)+
- 662 mRNAs were isolated using poly(A)+ Dynabeads (Invitrogen). The sequencing

663 library was generated and sequenced, and the DMS modifications were computed as

- 664 previously described (Rouskin et al.).
- 665

666 mRNA secondary structure prediction

667 Near the Ire1 cleavage sites, we first identified the most highly reactive base and set

668 its DMS modification signal as 1. Then, the DMS modification signal raw data for

other bases was normalized proportionally. Finally, we put a 38-base-pair RNA

670 sequence (19 base pair upstream and downstream from the Ire1 cleavage site) into

671 the RNA secondary structure prediction program mfold (Zuker, 2003). Bases with

- 672 normalized DMS modification signals >0.2 were forced to be single stranded to
- 673 constrain the RNA folding prediction.
- 674
- 675

- **Reference**

678	ARAGON, T., VAN ANKEN, E., PINCUS, D., SERAFIMOVA, I. M., KORENNYKH, A. V.,			
679	RUBIO, C. A. & WALTER, P. 2009. Messenger RNA targeting to endoplasmic			
680	reticulum stress signalling sites. <i>Nature</i> , 457, 736-40.			
681				
682	BERTOLOTTI, A., WANG, X., NOVOA, I., JUNGREIS, R., SCHLESSINGER, K., CHO, J. H.,			
683	WEST, A. B. & RON, D. 2001. Increased sensitivity to dextran sodium sulfate			
684	colitis in IRE1beta-deficient mice. J Clin Invest, 107, 585-93.			
685				
686	CALFON, M., ZENG, H., URANO, F., TILL, J. H., HUBBARD, S. R., HARDING, H. P.,			
687	CLARK, S. G. & RON, D. 2002. IRE1 couples endoplasmic reticulum load to			
688	secretory capacity by processing the XBP-1 mRNA. <i>Nature</i> , 415, 92-6.			
689				
690	COX, J. S., SHAMU, C. E. & WALTER, P. 1993. Transcriptional induction of genes			
691	encoding endoplasmic reticulum resident proteins requires a			
692	transmembrane protein kinase. <i>Cell</i> , 73, 1197-206.			
693				
694	CREDLE, J. J., FINER-MOORE, J. S., PAPA, F. R., STROUD, R. M. & WALTER, P. 2005. On			
695	the mechanism of sensing unfolded protein in the endoplasmic reticulum.			
696	Proc Natl Acad Sci U S A. 102, 18773-84.			
697				
698	GARDNER, B. M. & WALTER, P. 2011, Unfolded proteins are Ire1-activating ligands			
699	that directly induce the unfolded protein response. <i>Science</i> , 333, 1891-4.			
700				
701	GONZALEZ, T. N., SIDRAUSKI, C., DORFLER, S. & WALTER, P. 1999, Mechanism of			
702	non-spliceosomal mRNA splicing in the unfolded protein response pathway.			
703	<i>EMBO I</i> . 18, 3119-32.			
704				
705	GUTHRIE, C. & FINK, G. R. 2002, Guide to yeast genetics and molecular and cell			
706	<i>biology</i> . Amsterdam : Boston : London. Academic Press.			
707				
708	GUYDOSH, N. R., KIMMIG, P., WALTER, P. & GREEN, R. 2017, Regulated Ire1-			
709	dependent mRNA decay requires no-go mRNA degradation to maintain			
710	endoplasmic reticulum homeostasis in S. pombe. <i>Elife</i> . 6. e29216			
711	······································			
712	HOLLIEN, I., LIN, I. H., LI, H., STEVENS, N., WALTER, P. & WEISSMAN, I. S. 2009.			
713	Regulated Ire1-dependent decay of messenger RNAs in mammalian cells. <i>I</i>			
714	Cell Biol. 186, 323-31.			
715				
716	HOLLIEN, I. & WEISSMAN, I. S. 2006. Decay of endoplasmic reticulum-localized			
717	mRNAs during the unfolded protein response <i>Science</i> 313, 104-7			
718				

719 720	HOOKS, K. B. & GRIFFITHS-JONES, S. 2011. Conserved RNA structures in the non- canonical Hac1/Xbp1 intron. <i>RNA Biol</i> , 8, 552-6.
721 722 723 724	IMAGAWA, Y., HOSODA, A., SASAKA, S., TSURU, A. & KOHNO, K. 2008. RNase domains determine the functional difference between IRE1alpha and IRE1beta. <i>FEBS Lett,</i> 582, 656-60.
725 726 727 728 729	IWAWAKI, T., HOSODA, A., OKUDA, T., KAMIGORI, Y., NOMURA-FURUWATARI, C., KIMATA, Y., TSURU, A. & KOHNO, K. 2001. Translational control by the ER transmembrane kinase/ribonuclease IRE1 under ER stress. <i>Nat Cell Biol</i> , 3, 158-64.
730 731 732 733 734 735	JURKIN, J., HENKEL, T., NIELSEN, A. F., MINNICH, M., POPOW, J., KAUFMANN, T., HEINDL, K., HOFFMANN, T., BUSSLINGER, M. & MARTINEZ, J. 2014. The mammalian tRNA ligase complex mediates splicing of XBP1 mRNA and controls antibody secretion in plasma cells. <i>EMBO J</i> , 33, 2922-36.
735 736 737 738 720	KARAGOZ, G. E., ACOSTA-ALVEAR, D., NGUYEN, H. T., LEE, C. P., CHU, F. & WALTER, P. 2017. An unfolded protein-induced conformational switch activates mammalian IRE1. <i>Elife,</i> 6, e30700
739 740 741 742 742	KIMMIG, P., DIAZ, M., ZHENG, J., WILLIAMS, C. C., LANG, A., ARAGON, T., LI, H. & WALTER, P. 2012. The unfolded protein response in fission yeast modulates stability of select mRNAs to maintain protein homeostasis. <i>Elife</i> , 1, e00048.
743 744 745 746	KOHRER, K. & DOMDEY, H. 1991. Preparation of high molecular weight RNA. <i>Methods Enzymol</i> , 194, 398-405.
747 748 749 750	KORENNYKH, A. V., EGEA, P. F., KOROSTELEV, A. A., FINER-MOORE, J., ZHANG, C., SHOKAT, K. M., STROUD, R. M. & WALTER, P. 2009. The unfolded protein response signals through high-order assembly of Ire1. <i>Nature</i> , 457, 687-93.
751 752 753 754 755	KOSMACZEWSKI, S. G., EDWARDS, T. J., HAN, S. M., ECKWAHL, M. J., MEYER, B. I., PEACH, S., HESSELBERTH, J. R., WOLIN, S. L. & HAMMARLUND, M. 2014. The RtcB RNA ligase is an essential component of the metazoan unfolded protein response. <i>EMBO Rep</i> , 15, 1278-85.
756 757 759	LU, Y., LIANG, F. X. & WANG, X. 2014. A synthetic biology approach identifies the mammalian UPR RNA ligase RtcB. <i>Mol Cell</i> , 55, 758-70.
758 759 760	MAUREL, M., CHEVET, E., TAVERNIER, J. & GERLO, S. 2014. Getting RIDD of RNA: IRE1 in cell fate regulation. <i>Trends Biochem Sci</i> , 39, 245-54.
762 763 764	MOORE, K. & HOLLIEN, J. 2015. Ire1-mediated decay in mammalian cells relies on mRNA sequence, structure, and translational status. <i>Mol Biol Cell</i> , 26, 2873-84.

765	
766	MORENO S KLAR A & NURSE P 1991 Molecular genetic analysis of fission yeast
767	Schizosaccharomyces nombe. <i>Methods Enzymol.</i> 194, 795-823.
768	
769	MORI K MA W GETHING M I & SAMBROOK I 1993 A transmembrane protein
770	with a cdc^2+/CDC^28 -related kinase activity is required for signaling from the
771	FR to the nucleus <i>Cell</i> 74, 743-56
772	
773	NAKAMUDA D TSUDU A IKECAMI K IMACAWA Y EUUMOTO N & KOHNO K
774	2011 Mammalian EP stross consor IDE1 hota specifically down-regulates the
775	supposes of socratory nathway protains FERS Latt 585 122-8
776	synthesis of secretory pathway proteins. <i>PEDS Lett</i> , 303, 133-0.
770	NIWA M DATH C K DEDISI I & WAITED D 2005 Conomo scalo approachos for
770	discovering nevel nonconventional calicing substrates of the Iro1 nuclease
770	Conomo Biol 6, D2
780	Genome Dioi, 0, KS.
700	DESCHER I ACOSTA-ALVEAD D MENDEZ A S & WALTED D 2015 A
701	conformational BNA zinner promotes intron ejection during non-
702	conventional XRD1 mDNA colliging EMPO Page 16, 1609,00
703	conventional ADF1 mKWA splicing. EMDO Rep, 10, 1000-90.
704	DI LIMP D 7HANC 7 D ADDATHIDALS & MADIADDAN M 2015 A functional link
705	PLOMD, N., ZHANG, Z. N., APPATHUNAI, S. & MANIAPPAN, M. 2015. A functional fills
700	Elife A o07426
707	<i>Elije</i> , 4, e07420.
700	POND & WAITED D 2007 Signal integration in the endeplacmic reticulum
709	unfolded protein response. Nat Day Mol Call Piol 9, 510, 20
790	uniolaeu protein response. <i>Nat Kev Mol Cell Diol,</i> 6, 519-29.
791	ROUSKIN S ZUBRADT M WASHIETI S KELLIS M & WEISSMAN I S 2014
702	Conomo-wido probing of PNA structure reveals active unfolding of mPNA
793	structures in vivo. Natura 505, 701, 5
794	sti uctures in vivo. <i>Nuture</i> , 503, 701-5.
795	DIIRIO C DINCUS D VODENNVVH A SCHUCK S EI SAMAD H & WAITED D
790	2011 Homoostatic adaptation to ondenlasmic raticulum stress depends on
700	Irol kinaco activity I Call Piol 102, 171, 94
790	11 e1 Killase activity. J cell Diol, 195, 17 1-04.
000	SIDDAUSKI C. COV I. S. & WALTED D. 1006 tDNA ligage is required for regulated
000	mDNA colliging in the unfolded protein response. Call 97, 405, 12
001	mixing splicing in the unfolded protein response. <i>Cell</i> , 87, 403-13.
002	TAM A D KOONG A C & NIWA M 2014 Incl has distinct establishing machanisms for
003	NDD1/UAC1 and sing and DDD. Call Day 0.050.0
004 005	λθε 1/ ΠΑC1 Splitting and KIDD. <i>Cell Kep, 9</i> , 850-8.
005	TCUDU A EUUMOTO N TAKAUACUI C CAITO M NAKAMUDA D UMANO M
000	ISUNU, A., FUJIMUTU, N., TANAHASHI, S., SAITU, M., NANAMUKA, D., IWANU, M., IMAMAKI T. KADOKUDA H. DON D. & KOUNO K. 2012. Nogeting foodbook
007	IVVAVVARI, I., RADURURA, I., RUN, D. & RUIINU, R. 2013. Negative feedback
000 000	by IKE 1 Deta optimizes much production in gobiet cells. <i>Proc Natl Acad Sci U</i>
009	JA, 110, 2004-7.
910 8	

811 812 813 814 815 816	VAN ANKEN, E., PINCUS, D., COYLE, S., ARAGON, T., OSMAN, C., LARI, F., GOMEZ PUERTA, S., KORENNYKH, A. V. & WALTER, P. 2014. Specificity in endoplasmic reticulum-stress signaling in yeast entails a step-wise engagement of HAC1 mRNA to clusters of the stress sensor Ire1. <i>Elife</i> , 3 , e05031.
010	WALTED D & DON D 2011 The unfolded protein regranges from stress nothway to
01/	homoostatia nogulation. Science 224, 1001 6
818	nomeostatic regulation. <i>Science</i> , 334, 1081-6.
819	
820	YANAGITANI, K., IMAGAWA, Y., IWAWAKI, T., HOSODA, A., SAITO, M., KIMATA, Y. &
821	KOHNO, K. 2009. Cotranslational targeting of XBP1 protein to the membrane
822	promotes cytopiasmic splicing of its own mRNA. <i>Mol Cell</i> , 34, 191-200.
823	
824	YANAGITANI, K., KIMATA, Y., KADOKURA, H. & KOHNO, K. 2011. Translational
825	pausing ensures membrane targeting and cytoplasmic splicing of XBP1u
826	mRNA. <i>Science,</i> 331, 586-9.
827	
828	YOSHIDA, H., MATSUI, T., YAMAMOTO, A., OKADA, T. & MORI, K. 2001. XBP1 mRNA
829	is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a
830	highly active transcription factor. <i>Cell</i> , 107, 881-91.
831	
832	ZUKER, M. 2003. Mfold web server for nucleic acid folding and hybridization
833	prediction. Nucleic Acids Res, 31, 3406-15.
834	
835	ZULETA, I. A., ARANDA-DIAZ, A., LI, H. & EL-SAMAD, H. 2014. Dynamic
836	characterization of growth and gene expression using high-throughput
837	automated flow cytometry. <i>Nat Methods</i> , 11, 443-8.
838	
839	
840	

Table 1:

Yeast strains used in this study. All strains are derived from WL001 and WL002. All Ire1 constructs listed below are 3x FLAG-tagged within their lumenal domains.

strain	species	description
yWL001	Sc	WT, mat A, <i>leu2-3,112 TRP1 can1-100 ura3-1 ADE2 his3-11,15</i>
yWL002	Sc	ire1∆::NAT ^R
yWL003	Sc	ire1∆::NAT ^R , HO::Sp IRE1
yWL004	Sc	<i>ire1∆::NAT^R, HO::Sp</i> Lum <i>Sc</i> Cyto <i>IRE1</i>
yWL005	Sc	<i>ire1∆::NAT^R, HO::Sc</i> Lum <i>Sp</i> Cyto <i>IRE1</i>
yWL006	Sc	ire1∆::NAT ^R , HO::Sc IRE1
yWL007	Sc	<i>ire1D</i> :: <i>NAT</i> ^{<i>R</i>} , <i>HO</i> :: <i>Sc</i> Lum/transmembrane/linker <i>Sp</i> KR <i>IRE1</i>
yWL008	Sc	WT, leu2::5'hac1-gfp-3'hac1
yWL009	Sc	ire1∆::NAT ^R , leu2::5'hac1-gfp-3'hac1
yWL010	Sc	ire1∆::NAT [®] , leu2::5'hac1-gfp-3'hac1, HO::Sp IRE1
yWL011	Sc	ire1Δ::NAT ^R , leu2::5'hac1-gfp-3'hac1, HO:: SpLumScCyto IRE1
yWL012	Sc	ire1Δ::NAT ^R , leu2::5'hac1-gfp-3'hac1, HO:: ScLumSpCyto IRE1
yWL013	Sc	ire1∆::NAT ^R , leu2::5'hac1-gfp-3'hac1, HO::Sc IRE1
yWL014	Sc	leu2::5'hac1-gfp-3'hac1, his3::pTdh3-mCherry
yWL015	Sp	WT, mat h+, <i>ade6-M210, ura4-D18, leu1-32</i>
yWL016	Sp	ire1∆::KAN ^R
yWL017	Sp	ire1∆::KAN ^R , leu1::Sp IRE1
yWL018	Sp	ire1Δ::KAN ^R , leu1::SpLumScCyto IRE1
yWL019	Sp	ire1 <i>Δ::KAN^R, leu1::Sc</i> Lum <i>Sp</i> Cyto <i>IRE1</i>
yWL020	Sp	ire1Δ::KAN ^R , leu1::Sc IRE1
yWL021	Sp	ire1Δ::KAN ^R , leu1::SpLumScCyto IRE1, ura4::5'hac1-gfp-3'hac1
yWL022	Sp	ire1 <i>\Delta::KAN^R</i> , leu1::ScLumSpCyto IRE1, ura4::5'hac1-gfp-3'hac1
yWL023	Sp	ire1∆::KAN ^R , leu1::Sc IRE1, ura4::5'hac1-gfp-3'hac1
yWL024	Sp	<i>bip1::bip1-hac1</i> hybrid
yWL025	Sp	<i>bip1::bip1-hac1</i> hybrid A->U
yWL026	Sp	<i>bip1::bip1</i> splicing variant
yWL027	Sp	ura4::synthetic splicing substrate

Table 1:

849 qPCR primers used in this study.

qPCR primers description	sequence
uncleaved Sp BiP1 mRNA forward primer	GAATCGTGACTCTATAGCCATTAACA
uncleaved Sp BiP1 mRNA reverse primer	CAATTATTGTCAGTTCCACAAAGC
total Sp BiP1 mRNA forward primer	TGGTAAGGTTGATCCCGAAG
total <i>Sp BiP1</i> mRNA reverse primer	CATCGAGTTTTTGACGCTGA
Sp GAS2 mRNA forward primer	GTTGTCAACAATGCCTCGAA
<i>Sp GAS2</i> mRNA reverse primer	CGGTCTCAGAGTTGGTGTCA
Sp NDA2 mRNA forward primer	TCCATGAATCCAACAGCGTA
Sp NDA2 mRNA reverse primer	CTAGTAACGGCAGCCTGGAC













lane: 1 2 3 4







AA

bar: 1 2 3 4 5 6

Fig. 3-figure supplement 1 Li et al.

А

systematic ID	gono nomo	Iral alaguaga sita saguanga
systematic ID	gene name	ner cleavage site sequence
SPBC29A10.08	GAS2	ACGAUCCCUGUCGGUUAUGCUGGUGCCGAUAUUCCCGU
SPAC22A12.15c	BIP1	AA C ACCUAGUUAACUGG <mark>UGC</mark> UUUGUUAUCUUUGUAUUG
SPAC26A3.01	SXA1	AUUAUCGGUUUGUCAGU <mark>UGC C</mark> AUGACUAUUACUGGUAU
SPBC25H2.06c	HRF1	GAGCUAUUUGGCCUUCG <mark>UGC</mark> UAGUAAGGCUUGUGCUGU
SPAC1A6.04c	PLB1	UCCUUUGUGGCCUUUGU <mark>UGC</mark> A A AAGGGUCGUGAUGUCG
SPBP4G3.02	PHO1	AUUUGCAGUUAUGAAAU <mark>UGC</mark> CCUUCA A GACUAU A GCGA
SPCC970.03		CAUUU A UCGUUUCCGUC <mark>UGC</mark> CUCGCGCU AA CGAUGUUC
SPBC29A10.08	GAS2	GCUGU C GCUU A CGUUCG <mark>UGC</mark> UGCCGUUCGUGAUUCCAA
SPAC1A6.04c	PLB1	GUUGCCGA AAA GGCC AAUGC UGGCUUU AACA UC A GUCU
SPBP4G3.02	PHO1	AUCUUUGGAGGUGCCUA <mark>UGC</mark> UA A UAGCCUUGCAAAUUC
SPBC3D6.02	BUT2	CUU A GUGCUGACAUUCC <mark>UGC</mark> CAAGUCUAGCCGU C UCUU
SPAC56F8.07		CUUUUACUACGACAUGG <mark>UGC</mark> UGCAUGUUCGAGUUAUUU
SPCC830.08c	YOP1	GCCUUCUUUAGUAUCAA <mark>UGC</mark> U A UUGA AAC UACUA AC AA
SPAC4G9.15		GGAAUCGGUAAAGAGUAUGC UACUCAAUUAGCCAUGUC

YOP1

В



SPAC4G9.15

Fig 4 Li et al.

5' UTR | mutant |

ORF |

I.

ī.





2

T

with

splicing cassette

Fig. 4-figure supplement 1 Li et al.





A

IRE1alpha	1	LEKQLQFFQDVSDRIEKESLDGPIVKQLERGGRAVVKMDWRENITVPLQTDLRKFRTYKG
IRE1beta	1	RAKOLOFFODVSDWLEKESEOEPLVRALEAGGCAVVRDNWHEHISMPLOTDLRKFRSYKG
Sc_IRE1	1	KSKKLEFLLKVSDRIEIENRDPPSALIMKFDAGSDFV PSGDWTVKFDK FMDNLERVRK
Sp_IRE1	1	YAKKIDELIDVSDRFEVEERDPPSPILQMLENNSKSVIGENWTTCLHSSLVDNLGKYRKY
IRE1alpha	61	GSVRDLLRAMRNKKHHYRELPAEVRETLGSLPDDFVCYFTSRFPHLLAHTYRAMELCSHE
IRE1beta	61	TSVRDLLRAVRNKKHHY <mark>R</mark> ELPVEVRQALGQVPDGFVQYFT <mark>NRF</mark> PRLLLHTHRAMRSCASE
Sc_IRE1	61	YHSSKIMDLLRALRNKYHHFMDLPEDIAEIMGPVPDGFYDYFTKRFPNLLIGVYMIVKEN
Sp_IRE1	61	DGSKILDILRVLRNKRHHYQDLPESVRRVLGDLPDGFTSYFVEKFPMLLLHCYHLVKDVL
IRE1alpha	121	RLFQPVYFHEPPEPQPPVTPDAL
IRE1beta	121	SLFLPYYPPDSEARRPCPGATGR
Sc_IRE1	121	LSDDQILREFLYS
Sp_IRE1	121	YEESQFKRYLEV

